

Explicitly Computing With Modular Abelian Varieties

William Stein
Harvard University

February 6, 2004

Center for Communications Research in Princeton

```
J := J0(65); J;  
Modular abelian variety J0(65) of dimension 2 and level 65  
D := Decomposition(J); D;  
[  
  Modular abelian variety 85A of dimension 1, level 65 and  
  conductor 5*13 over Q.  
  Modular abelian variety 85B of dimension 1, level 65 and  
  conductor 5^2*13^2 over Q.  
  Modular abelian variety 85C of dimension 1, level 65 and  
  conductor 5^2*13^2 over Q.  
]  
#(D[1] meet D[2]);  
IsIsomorphic(D[2], Dual(D[2]));  
true  
Homomorphism from 85B to modular abelian variety of  
dimension 2 given on integral homology by  
0 0 -1  
1 0 -1  
1 1 -1  
L-Series(D[2]);  
L-Series(D[2]);  
L-Series(D[2]);
```

Overview of Talk

1. Modular Abelian Varieties
2. Computing With Modular Abelian Varieties

Modular Abelian Varieties

Abelian variety: A complete group variety



Abel


Examples:

1. Elliptic curves, e.g., $y^2 = x^3 + ax + b$
2. Jacobians of curves
3. Quotients of Jacobians of curves

Connection with Cryptography

Modular abelian varieties over finite fields provide a large source of groups that can be used for cryptography (e.g., Elliptic Curve Cryptography). I will focus on modular abelian varieties over infinite fields today, but the results are relevant for understanding the reductions of those varieties modulo primes.

The Modular Curve $X_1(N)$




Hecke

Let $\mathfrak{h}^* = \{z \in \mathbf{C} : \Im(z) > 0\} \cup \mathbf{P}^1(\mathbf{Q})$.

1. $X_1(N)_{\mathbf{C}} = \Gamma_1(N) \backslash \mathfrak{h}^*$ (compact Riemann surface)
2. $X_1(N)$ has natural structure of algebraic curve over \mathbf{Q}
3. $X_1(N)(\mathbf{C}) = \{(E, P) : \text{ord}(P) = N\} / \sim$ (moduli space)

N	≤ 10	11	13	37	169	512	2003
genus($X_1(N)$)	0	1	2	40	1070	7809	166167

Modular Forms



Hecke

1. Cuspidal modular forms (of weight 2):

$$S_2(N) = H^0\left(X_1(N), \Omega_{X_1(N)}^1\right)$$

2. $f \in S_2(N)$ has Fourier expansion in terms of $q(z) = e^{2\pi iz}$

$$f = \sum_{n=1}^{\infty} a_n q^n$$

3. Hecke algebra (*commutative* ring):

$$\mathbf{T} = \mathbf{Z}[T_1, T_2, \dots] \hookrightarrow \text{End}(S_2(N))$$

The Modular Jacobian $J_1(N)$



Jacobi

1. Jacobian of $X_1(N)$:

$$J_1(N) = \text{Jac}(X_1(N))$$

2. $J_1(N)$ is an abelian variety over \mathbf{Q} of dimension $g(X_1(N))$.
3. The elements of $J_1(N)$ parameterize divisor classes on $X_1(N)$ of degree 0.

Modular Abelian Varieties

A **modular abelian variety** A over a number field K is any abelian variety A (over K) such that there is a homomorphism

$$A \rightarrow J_1(N)$$

with finite kernel.



Shimura

Examples and Conjectures

Suppose $\dim A = 1$.

- **Theorem (Wiles, Breuil, Conrad, Diamond, Taylor).**
If $K = \mathbb{Q}$ then A is modular.
- **Theorem (Shimura).** If A has CM then A is modular.
- **Definition:** A over $\overline{\mathbb{Q}}$ is a **\mathbb{Q} -curve** if for each Galois conjugate A^σ of A there is an isogeny $A \rightarrow A^\sigma$ (an isogeny is a map with finite kernel).
Conjecture (Ribet, Serre). Over $\overline{\mathbb{Q}}$ the non-CM modular elliptic curves are exactly the \mathbb{Q} -curves.

GL₂-type

Defn. A simple abelian variety A/\mathbf{Q} is of GL₂-type if


$$\text{End}_0(A/\mathbf{Q}) = \text{End}(A/\mathbf{Q}) \otimes \mathbf{Q}$$

is a number field of degree $\dim(A)$.

Shimura associated GL₂-type modular abelian varieties to \mathbf{T} -eigenforms:

$$f = q + \sum_{n \geq 2} a_n q^n \in S_2(N)$$

$$I_f = \text{Ker}(\mathbf{T} \rightarrow \mathbf{Q}(a_1, a_2, a_3, \dots)), \quad T_n \mapsto a_n$$



Ken Ribet

Abelian variety A_f over \mathbf{Q} of $\dim = [\mathbf{Q}(a_1, a_2, \dots) : \mathbf{Q}]$:

$$A_f := J_1(N)/I_f J_1(N)$$

Theorem (Ribet). Shimura's A_f is \mathbf{Q} -isogeny simple since

$$\text{End}_0(A_f/\mathbf{Q}) = \mathbf{Q}(a_2, a_3, \dots).$$

Also there is an isogeny $J_1(N) \sim \prod_f A_f$, where the product is over Galois-conjugacy classes of f .

Conjecture. (Ribet)

The simple modular abelian varieties A over \mathbf{Q} are exactly the simple abelian varieties over \mathbf{Q} of GL_2 -type.

Ribet proved that his conjecture follows from Serre's unproven conjectures on modularity of odd mod p Galois representations.

2. Computing With Abelian Varieties



Goal: Develop a systematic theory for computing with modular abelian varieties.

Basic Problems: Presentation, isogeny testing, isomorphism testing, endomorphism ring, enumeration.

Arithmetic Problems: Special values of L -functions, computing Shafarevich-Tate groups, Tamagawa numbers, enumerating elements of isogeny class.

Presentation

Modular abelian varieties can be specified in many ways:

- Equations
- Built from newform abelian varieties A_f
- Arise theoretically (e.g., Jacobians of Shimura curves).

For all our questions today we will view a modular abelian variety as being defined in the following way. Any modular abelian variety B can be obtained by quotienting an abelian subvariety $A \subset J_1(N)$ by a finite subgroup G . Thus we represent B by giving a pair (A, G) , where $G \subset A \subset J_1(N)$.

Specifying A

An inclusion $\varphi : A \hookrightarrow J_1(N)$ induces an inclusion on homology

$$H_1(A, \mathbf{Q}) \hookrightarrow H_1(J_1(N), \mathbf{Q}),$$

and A is completely determined by the image of $H_1(A, \mathbf{Q})$ in the vector space $H_1(J_1(N), \mathbf{Q})$.

We give A by giving a subspace $V = V_{\mathbf{Q}} \subset H_1(J_1(N), \mathbf{Q})$.

Specifying G

By the Abel-Jacobi theory there is a canonical isomorphism

$$J_1(N)(\mathbf{C}) \cong H_1(J_1(N), \mathbf{R}) / H_1(J_1(N), \mathbf{Z}).$$

Likewise $A(\mathbf{C}) \cong V_{\mathbf{R}} / V_{\mathbf{Z}}$, where $V_{\mathbf{Z}} = V \cap H_1(J_1(N), \mathbf{Z})$, so

$$A(\mathbf{C})_{\text{tor}} \cong V_{\mathbf{Q}} / V_{\mathbf{Z}}.$$

We give G by giving finitely many elements of $V_{\mathbf{Q}} / V_{\mathbf{Z}}$.

Recognition Problem

Problem: When does a subspace $V \subset H_1(J_1(N), \mathbb{Q})$ correspond to an abelian subvariety A of $J_1(N)$ over K ?

Solution: Given an isogeny decomposition of $J_1(N)$ over K as a direct sum of simple abelian varieties, I have an algorithm to solve this problem. (It is straightforward to compute such a decomposition when $K = \mathbb{Q}$.)

Problem: Given a group G defined by a finite list of elements of $V_{\mathbb{Q}}/V_{\mathbb{Z}}$, find the smallest number field over which G is defined. This is important because if G is defined over K , then $B = A/G$ is defined over K .

Solution??: I have not solved this problem, which is likely very difficult.

Modular Symbols



Manin

Modular symbols provide a presentation of

$$H_1(X_1(N), \mathbf{Z})$$

on which one can give formulas for Hecke and other operators. They have been intensively studied by Birch, Manin, Shokurov, Mazur, Merel, Cremona, and others.

```
> M := CuspidalSubspace(ModularSymbols(Gamma1(11)));  
> Basis(M);  
[  
  -1/5*{-1/2, 0} + -2/5*{-1/4, 0} + 3/5*{-1/7, 0} + -1/5*{7/15, 1/2}  
  -2/5*{-1/2, 0} + 1/5*{-1/4, 0} + 1/5*{-1/7, 0} + -2/5*{7/15, 1/2}  
]
```


Enumeration Problem Over \mathbb{Q}

Problem: Give an algorithm to systematically enumerate every modular abelian variety over \mathbb{Q} .

The isogeny classes of simple modular abelian varieties over \mathbb{Q} are in bijection with *newforms*, which are eigenvectors for Hecke operators in the space $S_2(\Gamma_1(N))$ of modular forms. Using the Atkin-Lehner-Li theory of newforms, modular symbols, and linear algebra, we can thus enumerate the isogeny classes over \mathbb{Q} .

I **do not know** how to find all abelian varieties in an isogeny class, except when A has dimension 1, where it is solved. Maybe at least find several by intersecting $A \subset J_1(N)$ with other abelian varieties over \mathbb{Q} , quotienting out by intersection, and proving quotient is not isomorphic to A .

Example

```
> Factorization(J1(17));
[*
<Modular abelian variety 17A of dimension 1, level 17
and conductor 17 over Q, [
Homomorphism from 17A to J1(17) given on integral
homology by:
[-3  1  2 -2  0 -2  2 -1  2  4]
[-2 -2  0  0  0  0  0  2  4  0]
]>,
<Modular abelian variety 17A[2] of dimension 4, level 17
and conductor 17^4 over Q, [
Homomorphism from 17A[2] to J1(17) (not printing
8x10 matrix)
]>
*]
```

Enumeration Problem Over $\overline{\mathbb{Q}}$

Problem: Give an algorithm to systematically enumerate every modular abelian variety over $\overline{\mathbb{Q}}$.

There is a huge amount of work by Shimura, Ribet, González, Lario, and others, but still nobody has given an algorithm to enumerate all isogeny classes of modular abelian varieties over $\overline{\mathbb{Q}}$ explicitly. By explicit, I mean in the sense of giving defining data, i.e., a pair $(V, G \subset V_{\mathbb{Q}}/V_{\mathbb{Z}})$.

Obstructions:

- Difficulty of constructing $\text{End}(A_f/\overline{\mathbb{Q}})$ explicitly (I have an algorithm, but it is *way too slow* to be useful)
- Difficulty of decomposing $A_f/\overline{\mathbb{Q}}$ as a product of simples, even given $\text{End}(A_f/\overline{\mathbb{Q}})$. Need a good “Meataxe” over \mathbb{Q} .

Computing Endomorphism Rings

Problem: Given a modular abelian variety A over K , compute $\text{End}(A)$ explicitly, i.e., give matrices in $\text{End}(V)$ that generate $\text{End}(A)$ as an abelian group.

Solution: When $A \subset J_1(N)$ is simple, $\text{End}(A) \otimes \mathbf{Q}$ is a skew field, which can be computed. For example, if $K = \mathbf{Q}$, then $A = A_f$ is attached to a newform and $\text{End}(A) \otimes \mathbf{Q}$ is generated by the image of the Hecke algebra. We can then find $\text{End}(A)$ in $\text{End}(A) \otimes \mathbf{Q}$ as the \mathbf{Z} -submodule of elements that preserve the lattice $V_{\mathbf{Z}}$.

We can also explicitly compute $\text{Hom}(A, B)$ for any modular abelian varieties A and B , by writing A and B as simples, computing endomorphism algebras, and finding the \mathbf{Z} -module of homomorphisms that induce a map that fixes integral homology.

Example

```
> A := J0(33); A;
Modular abelian variety J0(33) of dimension 3 and level 3*11 over Q
> End(A);
Group of homomorphisms from J0(33) to J0(33)
> Basis(End(A));
[
  Homomorphism from J0(33) to J0(33) (not printing 6x6 matrix),
  Homomorphism from J0(33) to J0(33) (not printing 6x6 matrix),
  Homomorphism from J0(33) to J0(33) (not printing 6x6 matrix),
  Homomorphism from J0(33) to J0(33) (not printing 6x6 matrix),
  Homomorphism from J0(33) to J0(33) (not printing 6x6 matrix)
]
> Matrix(Basis(End(A))[2]);
[ 0  1  0  0  0 -1]
[ 0  1  0  0  0  0]
[ 0  1  0  0 -1  0]
[ 0  1 -1  1 -1  0]
[ 0  1 -1  0  0  0]
[-1  1  0  0  0  0]
```

Isogeny Testing

Problem: Given modular abelian varieties A and B , determine whether or not A is isogenous to B .

Determine whether A is isogenous to B is easy, since we may assume A and B are attached to newforms $\sum a_n q^n$ and $\sum b_n q^n$, and then A is isogenous to B if and only if the newforms are Galois conjugate.

Isomorphism Testing

Problem: Suppose A is isogenous to B . Decide whether A is isomorphic to B .

I do not know how to do this in general. Assume we have computed $\text{End}(A)$, $\text{End}(B)$, and $\text{Hom}(A, B)$ explicitly. Given a basis for $\text{Hom}(A, B)$, how do we know if some linear combination of that basis has determinant 1? It's not clear (to me).

If A and B are both simple and have commutative endomorphism ring, then I found an algorithm to decide whether A is isomorphic to B . This algorithm can be extended to abelian varieties that are products of such A , assuming the factors occur with multiplicity 1 (up to isogeny). However, I do not know in general how to decide whether $A \oplus A$ is isomorphic to $B \oplus B$, though I have a vague strategy that I think might work.

Algorithm for Testing Isomorphism

Suppose A and B are explicitly defined modular abelian varieties over \mathbf{Q} that are both isogenous to an abelian variety A_f . The following algorithm determine whether A is isomorphic to B .

Let $H = \text{Hom}(A, B)$. Both A and B are given explicitly by pairs (V, G_1) and (V, G_2) , so we can compute an isogeny $f : B \rightarrow A$. Let $H_f = \{\phi \circ f : \phi \in H\} \subset \text{End}(B)$. Note that A is isomorphic to B if and only if H_f contains an element of degree $\deg(f)$. Also note that H_f has finite index in $\text{End}(B)$.

By hypothesis $K = \text{End}(B) \otimes \mathbf{Q}$ is the field generated by the Fourier coefficients of f . The norm of an element of K is the positive square root of the degree of the corresponding homomorphism (see Milne in Cornell-Silverman, pg 126, Prop. 12.12).

Thus if $\deg(f)$ is not a perfect square, then there can be no element of B of degree $\deg(f)$, so A is not isomorphic to B . Thus suppose $\deg(f) = d^2$.

Typically there will be infinitely many element in \mathcal{O}_K of norm d , but there are only finitely many up to units. There is an algorithm, which involves computing the class group of \mathcal{O}_K , which enumerates representative elements of \mathcal{O}_K of norm d , up to units (e.g., the `NormEquation` command in MAGMA). Thus suppose we have computed representative elements z_1, \dots, z_n of the elements of \mathcal{O}_K with norm d . Then A is isomorphic to B if and only if there is a unit u and a z_i such that $u^{-1}z_i \in H_f \subset K$. Equivalently, such that $z_i \in uH_f$. There are only finitely many possibilities for uH_f , since H_f has finite index in \mathcal{O}_K and $[\mathcal{O}_K : uH_f] = [\mathcal{O}_K : H_f]$, since $\mathcal{O}_K = u\mathcal{O}_K$. We can thus list all subgroups uH_f (since we can compute generators for \mathcal{O}_K^*) and hence determine whether H_f contains an element of norm d , as required.

**Thank you for
inviting me!**



Acknowledgements:

Papers of Ken Ribet, Goro Shimura, and Joan Lario.
Conversations with John Cannon, Enrique Gonzalez, David Ko-
hel, Barry Mazur, Allan Steel