

Homework 7 for Math 581F

Due FRIDAY November 16, 2007

Each problem has equal weight, and parts of problems are worth the same amount as each other.

- For each of the following three fields, determining if there is an order of discriminant 20 contained in its ring of integers:

$$K = \mathbb{Q}(\sqrt{5}), \quad K = \mathbb{Q}(\sqrt[3]{2}), \quad \text{and } \dots$$

K any extension of \mathbb{Q} of degree 2005. [Hint: for the last one, apply the exact form of our theorem about finiteness of class groups to the unit ideal to show that the discriminant of a degree 2005 field must be large.]

- Compute the class group of $\mathbb{Q}(\sqrt{-15})$ following a similar approach to the computation of the class group of $\mathbb{Q}(\sqrt{10})$ in the book. (Do not do this by typing 1 or 2 lines into Sage, but instead compute the Minkowski bound, etc.)
- Prove that the quantity $C_{r,s}$ in our theorem about finiteness of the class group can be taken to be $(\frac{4}{\pi})^s \frac{n!}{n^n}$, as follows (adapted from [?, pg. 19]): Let S be the set of elements $(x_1, \dots, x_n) \in \mathbb{R}^n$ such that

$$|x_1| + \dots + |x_r| + 2 \sum_{v=r+1}^{r+s} \sqrt{x_v^2 + x_{v+s}^2} \leq 1.$$

- Prove that S is convex and that $M = n^{-n}$, where

$$M = \max\{|x_1 \cdots x_r \cdot (x_{r+1}^2 + x_{(r+1)+s}^2) \cdots (x_{r+s}^2 + x_n^2)| : (x_1, \dots, x_n) \in S\}.$$

[Hint: For convexity, use the triangle inequality and that for $0 \leq \lambda \leq 1$, we have

$$\begin{aligned} \lambda \sqrt{x_1^2 + y_1^2} + (1 - \lambda) \sqrt{x_2^2 + y_2^2} \\ \geq \sqrt{(\lambda x_1 + (1 - \lambda)x_2)^2 + (\lambda y_1 + (1 - \lambda)y_2)^2} \end{aligned}$$

for $0 \leq \lambda \leq 1$. In polar coordinates this last inequality is

$$\lambda r_1 + (1 - \lambda)r_2 \geq \sqrt{\lambda^2 r_1^2 + 2\lambda(1 - \lambda)r_1 r_2 \cos(\theta_1 - \theta_2) + (1 - \lambda)^2 r_2^2},$$

which is trivial. That $M \leq n^{-n}$ follows from the inequality between the arithmetic and geometric means.

- Transforming pairs x_v, x_{v+s} from Cartesian to polar coordinates, show also that $v = 2^r (2\pi)^s D_{r,s}(1)$, where

$$D_{\ell,m}(t) = \int \cdots \int_{\mathcal{R}_{\ell,m}(t)} y_1 \cdots y_m dx_1 \cdots dx_\ell dy_1 \cdots dy_m$$

and $\mathcal{R}_{\ell,\uparrow}(t)$ is given by $x_\rho \geq 0$ ($1 \leq \rho \leq \ell$), $y_\rho \geq 0$ ($1 \leq \rho \leq m$) and

$$x_1 + \cdots + x_\ell + 2(y_1 + \cdots + y_m) \leq t.$$

(c) Prove that

$$D_{\ell,m}(t) = \int_0^t D_{\ell-1,m}(t-x)dx = \int_0^{t/2} D_{\ell,m-1}(t-2y)ydy$$

and deduce by induction that

$$D_{\ell,m}(t) = \frac{4^{-m}t^{\ell+2m}}{(\ell+2m)!}$$