

Math 124 Problem Set 5

2. We establish both identities by induction.

Claim: $[a_n, a_{n-1}, \dots, a_1, a_0] = \frac{p_n}{p_{n-1}}$.

Since $p_{-1} = 1$ and $p_0 = a_0$, $[a_0] = \frac{p_0}{p_{-1}}$. This establishes the base case. Now suppose that $[a_{n-1}, \dots, a_0] = \frac{p_{n-1}}{p_{n-2}}$. Then $[a_n, a_{n-1}, \dots, a_0] = a_n + 1/[a_{n-1}, \dots, a_0] = a_n + \frac{1}{\frac{p_{n-1}}{p_{n-2}}} = a_n + \frac{p_{n-2}}{p_{n-1}} = \frac{a_n p_{n-1} + p_{n-2}}{p_{n-1}}$. The numerator is just the definition of p_n ; this proves the claim.

Claim: $[a_n, a_{n-1}, \dots, a_1] = \frac{q_n}{q_{n-1}}$.

Since $q_0 = 1$ and $q_1 = a_1$, $[a_1] = \frac{q_1}{q_0}$. This establishes the base case. Now suppose that $[a_{n-1}, \dots, a_1] = \frac{q_{n-1}}{q_{n-2}}$. Then $[a_n, a_{n-1}, \dots, a_1] = a_n + 1/[a_{n-1}, \dots, a_1] = a_n + \frac{1}{\frac{q_{n-1}}{q_{n-2}}} = a_n + \frac{q_{n-2}}{q_{n-1}} = \frac{a_n q_{n-1} + q_{n-2}}{q_{n-1}}$. The numerator is just the definition of q_n ; this proves the claim.

3. If we compute $ellj(t)$ in PARI, where $t = -0.5 + 0.3281996289i$ the result is $61.7142856 \dots - 6.2E-26I$. The imaginary part is effectively 0, so we wish to guess the rational number that gives the real part. The command $contfrac(61.7142856)$ gives $[61, 1, 2, 2, 178571]$, suggesting that a good guess for our rational number is given by the continued fraction $[61, 1, 2, 2]$. This value is $\frac{432}{7}$.

4i. Let $\alpha = [2, 3]$. Then $\alpha = 2 + \frac{1}{3 + \frac{1}{\alpha}}$, so $3\alpha^2 - 6\alpha - 2 = 0$. Solving for α yields $1 + \frac{\sqrt{13}}{3}$.

4ii. First we compute $\alpha = [1, 2, 1]$. This gives $\alpha = 1 + (2 + \frac{1}{(1 + \alpha^{-1})})^{-1}$. Solving for α yields $3\alpha^2 - 2\alpha - 3 = 0$, so $\alpha = \frac{1 + \sqrt{10}}{3}$. Now $[2, \overline{1, 2, 1}] = 2 + \frac{1}{[1, 2, 1]}$, so our final answer is $\frac{5 + \sqrt{10}}{3}$.

4iii. This is $[\overline{1, 2, 3}]^{-1}$. As above, if $\alpha = [1, 2, 3]$ then $\alpha = 1 + (2 + \frac{1}{3 + \frac{1}{\alpha}})^{-1}$. This simplifies to $\alpha = 7\alpha^2 + 8\alpha - 3 = 0$, so $\alpha = \frac{4 + \sqrt{37}}{7}$. Therefore our desired answer is $\frac{-4 + \sqrt{37}}{3}$.

5. For all three parts we use $contfrac$ in PARI to find the continued fraction and prove that the answer is correct.

5i. We claim that $\sqrt{5} = [2, \overline{4}]$. Let $\alpha = [\overline{4}]$; then $\alpha = 4 + \frac{1}{\alpha}$, so $\alpha = 2 + \sqrt{5}$. Now $[2, \overline{4}] = 2 + \frac{1}{2 + \sqrt{5}} = \sqrt{5}$, as desired.

5ii. We claim that $\frac{1 + \sqrt{13}}{2} = [2, \overline{3}]$. Let $\alpha = [\overline{3}]$; then $\alpha = 3 + \frac{1}{\alpha}$, so $\alpha^2 - 3\alpha - 1 = 0$. This gives $\alpha = \frac{3 + \sqrt{13}}{2}$. Then $[2, \overline{3}] = 2 + \frac{2}{3 + \sqrt{13}} = \frac{1 + \sqrt{13}}{2}$.

5iii. We claim that $\frac{5 + \sqrt{37}}{4} = [2, \overline{1, 3}]$. Let $\alpha = [2, \overline{1, 3}]$; then $\alpha = 2 + (1 + (3 + \frac{1}{\alpha})^{-1})^{-1}$, so $4\alpha^2 - 10\alpha - 3 = 0$. Solving for α gives $\frac{5 + \sqrt{37}}{4}$, as desired.

6i. First we compute $[\overline{2n}]$. Let $\alpha = [\overline{2n}]$; then $\alpha = 2n + \frac{1}{\alpha}$. This gives $\alpha^2 - 2n\alpha - 1 = 0$, so $\alpha = n + \sqrt{n^2 + 1}$. Now $[n, \overline{2n}] = n + \frac{1}{\alpha} = n - (n - \sqrt{n^2 + 1}) = \sqrt{n^2 + 1}$, as desired.

6ii. Using the previous part, we know that $\sqrt{5} = [2, \overline{4}]$. We can try successive convergents until two agree up to four decimal places; once such convergent is $682/305$.

7. In PARI, use $convergents(contfrac(\pi))$ to obtain the convergents of the continued fraction of π . We can now test convergents for property described in the problem, noting that smaller denominators are more likely to work. One convergent that satisfies the property is $3/1$, since $\pi - 3 < .15$ and $\frac{1}{\sqrt{5}} > .44$. The next is $22/7$, since $22/7 - \pi < .002$ and $\frac{1}{49\sqrt{5}} > .009$. A third is $355/113$, since $355/113 - \pi < .0000003$

while $\frac{1}{113^2\sqrt{5}} > .00003$.

8. In PARI, the command `contfrac(exp(2))` gives

[7, 2, 1, 1, 3, 18, 5, 1, 1, 6, 30, 8, 1, 1, 9, 42, 11, 1, 1, 12, 54, 14, 1, 1, 15, 77, 17, 1, 1, 18, 78, 20, 1, 1, 21, 90, ...]

This suggests that after the initial 7, the i th group of 5 numbers is of the form

$$3(i - 1) + 2, 1, 1, 3i, 18 + 12(i - 1).$$

9i. We are looking for natural number solutions to $n + 1 = x^2$, $\frac{n}{2} + 1 = y^2$. Isolating n yields $n = x^2 - 1 = 2(y^2 - 1)$, implying that $x^2 - 2y^2 = -1$. There are infinitely many solutions (x, y) to this equation if the period of the continued fraction of $\sqrt{2}$ has odd order. Indeed, $\sqrt{2} = [1, \overline{2}]$. For each (x, y) we can take $2(y^2 - 1)$ to find a unique (even) n ; thus there are infinitely many n satisfying the desired property.

9ii. We can use the convergents program introduced in class to produce a list of convergents for the continued fraction of `sqrt2`. The odd terms are of interest, and since we want to find $n > 389$, we want the denominator of the convergent to be at least 14. The first two such convergents are $\frac{41}{29}$ and $\frac{239}{169}$. They yield $n = 2(29^2 - 1) = \mathbf{1680}$ and $n = 2(169^2 - 1) = \mathbf{57120}$.

10. If x and y are consecutive integers, then we have $z^2 = x^2 + (x + 1)^2 = 2x^2 + 2x + 1$. Multiplying by 2, we have $2z^2 = 4x^2 + 4x + 2 = (2x + 1)^2 + 1$. Put $u = 2x + 1$; then $u^2 - 2z^2 = -1$. From the previous problem we know there are infinitely many solutions to this equation for u and z . Clearly each solution (u, z) gives a unique (x, z) . Lastly we verify that each of these solutions (x, z) leads to a primitive Pythagorean triple. But this is trivial: clearly x and $x + 1$ are coprime, and if either shares a nontrivial divisor with z then the third must also share this divisor, implying that $(x, x + 1) > 1$, which is impossible.