



Mean Values of Derivatives of Modular \mathbb{L} -Series

M. Ram Murty, V. Kumar Murty

The Annals of Mathematics, Second Series, Volume 133, Issue 3 (May, 1991), 447-475.

Your use of the JSTOR database indicates your acceptance of JSTOR's Terms and Conditions of Use. A copy of JSTOR's Terms and Conditions of Use is available at <http://www.jstor.org/about/terms.html>, by contacting JSTOR at jstor-info@umich.edu, or by calling JSTOR at (888)388-3574, (734)998-9101 or (FAX) (734)998-9113. No part of a JSTOR transmission may be copied, downloaded, stored, further transmitted, transferred, distributed, altered, or otherwise used, in any form or by any means, except: (1) one stored electronic and one paper copy of any article solely for your personal, non-commercial use, or (2) with prior written permission of JSTOR and the publisher of the article or other text.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

The Annals of Mathematics is published by The Annals of Mathematics. Please contact the publisher for further permissions regarding the use of this work. Publisher contact information may be obtained at <http://www.jstor.org/journals/annals.html>.

The Annals of Mathematics
©1991 The Annals of Mathematics

JSTOR and the JSTOR logo are trademarks of JSTOR, and are Registered in the U.S. Patent and Trademark Office. For more information on JSTOR contact jstor-info@umich.edu.

©2001 JSTOR

Mean values of derivatives of modular L -series

By M. RAM MURTY AND V. KUMAR MURTY

1. Introduction

Recently, Kolyvagin [4] proved the finiteness of the Tate-Shafarevic group $\text{III}_{E/\mathbf{Q}}$ of certain modular elliptic curves E over \mathbf{Q} . More precisely, let E/\mathbf{Q} be a modular elliptic curve with conductor N and $L(s)$ its associated L -series:

$$L(s) = \sum_{n=1}^{\infty} \frac{a(n)}{n^s}.$$

Set

$$L_D(s) = \sum_{n=1}^{\infty} \frac{a(n)}{n^s} \left(\frac{D}{n} \right)$$

where (D/n) is the Legendre symbol. Suppose that $L(1) \neq 0$ and there exists a $D < 0$, such that

(i) $L_D(s)$ has a simple zero at $s = 1$ and

(ii) all primes which divide the conductor of E split in the imaginary quadratic field $\mathbf{Q}(\sqrt{D})$.

Under these conditions, Kolyvagin [4] showed that both $E(\mathbf{Q})$ and $\text{III}_{E/\mathbf{Q}}$ are finite. More recently, he extended this theorem to show that if $L(s)$ has a simple zero at $s = 1$ and there is a $D < 0$ satisfying (ii) above and $L_D(1) \neq 0$, then $\text{rank } E(\mathbf{Q}) = 1$ and $\text{III}_{E/\mathbf{Q}}$ is finite. Previously, Rubin [11] established the finiteness of $\text{III}_{E/\mathbf{Q}}$ for CM elliptic curves for which $L(1) \neq 0$. The work of Rubin and Kolyvagin represents significant steps toward the resolution of the important conjecture that $\text{III}_{E/\mathbf{Q}}$ is finite.

The purpose of this paper is to establish the existence of a $D < 0$ such that $L_D(s)$ has a simple zero at $s = 1$ and all primes dividing the conductor of E split completely in the quadratic field $\mathbf{Q}(\sqrt{D})$. Thus, the result of Kolyvagin can be stated without any hypothesis on quadratic twists of $L(s)$. We prove our theorems by showing that the mean value of $L'_D(1)$ is non-zero. More precisely, we prove the following.

THEOREM 1. Suppose that $L(1) \neq 0$. Let

$$C = \frac{1}{2N} \sum_{n_1, n_2} \frac{a(n_1 n_2^2)}{n_1 n_2^2} \frac{\phi(n_2)}{n_2},$$

where n_1 ranges over positive integers with the property that $p \mid n_1$ implies $p \mid 4N$ and $(n_2, 4N) = 1$ and ϕ denotes Euler's function. Then, $C \neq 0$ and

$$\sum_{\substack{0 < -D \leq Y \\ D \equiv 1 \pmod{4N}}} L'_D(1) = CY(\log Y) + o(Y \log Y),$$

as $Y \rightarrow \infty$.

The theorem is, in reality, a theorem about the mean values of derivatives of L -series attached to modular forms.

To fix ideas, let $F(z)$ be a cusp form of weight 2 on $\Gamma_0(N)$ which is a normalized eigenform for the Hecke operators. Suppose that F is not a modular form on $\Gamma_0(M)$ for any proper divisor M of N , and write

$$F(z) = \sum_{n=1}^{\infty} a(n) e^{2\pi i n z}$$

for its Fourier expansion at the cusp $i\infty$. Let

$$L(s) = \sum_{n=1}^{\infty} \frac{a(n)}{n^s}$$

be the associated L -series which satisfies a functional equation:

$$A^s \Gamma(s) L(s) = w A^{2-s} \Gamma(2-s) L(2-s),$$

where $w = \pm 1$ and $A = \sqrt{N}/2\pi$. Let $\chi_D(n) = (D/n)$ be a real character mod D . Then, we can consider

$$L_D(s) = \sum_{n=1}^{\infty} \frac{a(n)}{n^s} \left(\frac{D}{n} \right).$$

It extends to an entire function of s , and if D is a fundamental discriminant prime to N , then $L_D(s)$ satisfies a functional equation:

$$(A|D|)^s \Gamma(s) L_D(s) = w \chi_D(-N) (A|D|)^{2-s} \Gamma(2-s) L_D(2-s).$$

THEOREM 2. Suppose that $L(1) \neq 0$. Let

$$C = \frac{1}{2N} \sum_{n_1, n_2} \frac{a(n_1 n_2^2)}{n_1 n_2^2} \frac{\phi(n_2)}{n_2},$$

where n_1 ranges over positive integers with the property that $p \mid n_1$ implies

$p \mid 4N$ and $(n_2, 4N) = 1$ and ϕ denotes Euler's function. Then $C \neq 0$ and

$$\sum_{\substack{0 < -D \leq Y \\ D \equiv 1 \pmod{4N}}} L'_D(1) = CY \log Y + o(Y \log Y),$$

as $Y \rightarrow \infty$.

COROLLARY. Suppose that $L(1) \neq 0$. Then, there are infinitely many fundamental discriminants $D < 0$ such that $L_D(s)$ has a simple zero at $s = 1$ and all primes dividing the conductor of E split completely in the imaginary quadratic field $\mathbf{Q}(\sqrt{D})$.

This corollary was also established by Bump, Friedberg and Hoffstein [1] utilising the automorphic theory of $\mathrm{GSp}(4)$.

Remark 1. The proof will show that we need not assume that $L(1) \neq 0$ but only that the root number of $L(s)$ is $+1$.

Remark 2. We know that

$$w = (-1)^{\mathrm{ord}_{s=1} L(E, s)}.$$

Therefore, the assumption that $w = +1$ and the congruence $D \equiv 1 \pmod{4N}$ imply that $w\chi_D(-N) = -1$ and so $L_D(E, s)$ has an odd order zero at $s = 1$.

Remark 3. To see that C is non-zero, we obtain that

$$\begin{aligned} & \sum_{n_1, n_2} \frac{a(n_1 n_2^2)}{(n_1 n_2^2)^s} \frac{\phi(n_2)}{n_2} \\ &= \left(\sum_{\substack{b=1 \\ p \nmid b \Rightarrow p \mid 4N}}^{\infty} \frac{a(b)}{b^s} \right) \prod_{p \nmid 4N} \left(1 + \frac{p-1}{p} \left(\frac{a(p^2)}{p^{2s}} + \frac{a(p^4)}{p^{4s}} + \cdots \right) \right). \end{aligned}$$

Now consider the Euler product above. For $p \nmid 4N$, let us write $a(p) = \alpha_p + \beta_p$ with $|\alpha_p| = |\beta_p| = p^{1/2}$. If from the factor at p we factor out $\Sigma a(p^{2\alpha})/p^{2\alpha s}$, we find (for even N)

$$\sum \frac{a(n_1 n_2^2)}{(n_1 n_2^2)^s} \frac{\phi(n_2)}{n_2} = \mathcal{P}(s) \left(\sum_{(n, 4N)=1} \frac{a(n^2)}{n^{2s}} \right) \prod_{p \mid 4N} \left(1 - \frac{a(p)}{p^s} \right)^{-1}$$

where

$$\mathcal{P}(s) = \prod_{p \nmid 4N} \left(1 + \frac{1}{p} \left[\left(1 - \frac{1}{p^{4s-2}} \right)^{-1} \left(1 - \frac{\alpha_p^2}{p^{2s}} \right) \right. \right. \\ \left. \left. \times \left(1 - \frac{\beta_p^2}{p^{2s}} \right) \left(1 - \frac{1}{p^{2s-1}} \right) - 1 \right] \right).$$

We observe that this product converges absolutely for $\operatorname{Re}(s) > 1/2$ and that none of the Euler factors vanishes at $s = 1$. Since $a(p) = 0, \pm 1$, for $p \mid 4N$,

$$\prod_{p \mid 4N} \left(1 - \frac{a(p)}{p} \right)^{-1} \neq 0.$$

If N is odd, a similar non-zero factor is obtained. Moreover, the Euler product of the series

$$\sum_{\substack{n=1 \\ (n, 4N)=1}}^{\infty} \frac{a(n^2)}{n^s}$$

differs from that of $L(s, \operatorname{Sym}^2)\zeta(2s-2)^{-1}$ at only a finite number of primes and at these primes, none of the Euler factors vanishes at $s = 2$. Thus,

$$C \neq 0,$$

since $L(2, \operatorname{Sym}^2) \neq 0$ (see for example [5] or [8]). Therefore, our theorem shows that there exists D such that (i) and (ii) hold.

Remark 4. Our proof produces an error term of $\mathbf{O}(Y(\log Y)^{1-\rho})$ for an explicit value of ρ .

Remark 5. Our method is applicable to holomorphic modular forms f of any weight $k \geq 2$ for $\Gamma_0(N)$.

Proof of the corollary. Suppose there are only a finite number of D_i 's, D_1, \dots, D_r , say, satisfying (i) and (ii). Set

$$\frac{1}{L(s)} = \sum_{n=1}^{\infty} \frac{\tilde{\mu}(n)}{n^s}.$$

Notice that $|\tilde{\mu}(d)| \leq \mathbf{d}(d)\sqrt{d}$ and $\tilde{\mu}(d) = 0$ if $p^3 \mid d$ for any prime p . (We write $\mathbf{d}(n)$ for the number of positive divisors of n .) Then, fixing an i and writing

$D = D_i \delta^2$, we have the relation

$$L_D(s) = L_{D_i}(s) \sum_{d|\delta^2} \frac{\tilde{\mu}(d)}{d^s} \left(\frac{D_i}{d} \right).$$

We deduce that

$$L'_D(1) = L'_{D_i}(1) \sum_{d|\delta^2} \frac{\tilde{\mu}(d)}{d} \left(\frac{D_i}{d} \right).$$

Thus,

$$\begin{aligned} \sum_{\substack{0 < -D \leq Y \\ D = D_i \delta^2}} L'_D(1) &\ll |L'_{D_i}(1)| \sum_{0 \leq \delta \leq \sqrt{Y}} \sum_{d|\delta^2} \frac{|\tilde{\mu}(d)|}{d} \\ &\ll \sum_{d \leq Y} \frac{|\tilde{\mu}(d)|}{d} \left(\sum_{\substack{\delta \leq \sqrt{Y} \\ \delta^2 \equiv 0 \pmod{d}}} 1 \right). \end{aligned}$$

We write $d = d_0 d_1^2$ with d_0, d_1 coprime and squarefree. Then, the inner sum is

$$\frac{\sqrt{Y}}{d_0 d_1} + \mathbf{O}(1).$$

Then our sum is

$$\ll \sum_{d_1 \leq Y} \frac{\mathbf{d}(d_1^2)}{d_1} \sum_{d_0 \leq Y/d_1^2} \frac{\mathbf{d}(d_0)}{\sqrt{d_0}} \left(\frac{\sqrt{Y}}{d_0 d_1} + \mathbf{O}(1) \right)$$

and we easily deduce that this is

$$\ll \sqrt{Y} \log Y.$$

Summing over i produces a contradiction.

There are heuristics that suggest, in fact, that there should be a positive proportion of such D 's. To approach the problem of getting an estimate for the *number* of such D 's, one should modify the kernel in our integrals to make it more sensitive to the counting problem.

We close this section by introducing some further notation. If D_0 is a fundamental discriminant which is coprime to N , then

$$L_{D_0}(1+s) = w \chi_{D_0}(-N) |D_0|^{-2s} A^{-2s} \frac{\Gamma(1-s)}{\Gamma(1+s)} L_{D_0}(1-s).$$

Any $D \equiv 1 \pmod{4}$ can be written as $D = D_0 r^2$, where D_0 is a fundamental

discriminant and

$$L_D(s) = L_{D_0}(s) \sum_{d|r^2} \frac{\tilde{\mu}(d)}{d^s} \left(\frac{D_0}{d} \right).$$

Let us set

$$\begin{aligned} \tilde{f}_Y(n, s; a) &= \sum_{\substack{0 < -D \leq Y \\ D \equiv a \pmod{4N}}} \left(\frac{D}{n} \right) |D|^s, \quad D \text{ unrestricted} \\ f_Y(n, s; a) &= \sum_{\substack{0 < -D_0 \leq Y \\ D_0 \equiv a \pmod{4N}}} \left(\frac{D_0}{n} \right) |D_0|^s, \quad D_0 \text{ fundamental} \\ \tilde{f}_Y(n; a) &= \tilde{f}_Y(n, 0; a), \quad f_Y(n; a) = f_Y(n, 0; a). \end{aligned}$$

We shall write $\mathbf{d}(n)$ for the number of positive divisors of n .

We stress that the naive approach to the proof of the main theorem works after a few technical details are surmounted. The reader interested in ignoring these details and desiring an intuitive description of the proof can find it in [9].

The next two sections establish the requisite lemmas to estimate the sums we will encounter.

2. Lemmas

Throughout this paper, we will adopt the convention that a natural number n is written as $n_1 n_2$, where n_1 has the property that any prime divisor of it is also a divisor of $2N$ and n_2 is coprime to $2N$. On certain occasions, the same convention applies when we write $m = m_1 m_2$.

LEMMA 1. For $(a, N) = 1$,

$$\sum'_{\substack{n \leq X \\ n = n_1 n_2}} \left| \sum_{\substack{0 < -h \leq Y \\ h \equiv a \pmod{N}}} \left(\frac{h}{n} \right) \right|^2 \ll (N^2 / \phi(N)) XY \log^2 X$$

where the implied constant is absolute and the sum over $n \leq X$ such that n_2 is not a perfect square.

Proof. This lemma is easily derived from the results of Jutila [3] (for the case $N = 1$) and Fainleib and Saparnijazov [2] (for the general case). They prove:

$$(1) \quad \sum'_{\substack{n \leq X \\ (n, 2N) = 1}} \left| \sum_{\substack{0 < -h \leq Y \\ h \equiv a \pmod{N}}} \left(\frac{h}{n} \right) \right|^2 \ll NXY \log^2 X,$$

where the prime on the first sum means that n is not a square. The sum in the lemma is seen to be

$$\sum_{n_1 \leq X} \sum'_{n_2 \leq X/n_1} \left| \sum_{\substack{0 < -h \leq Y \\ h \equiv a \pmod{N}}} \left(\frac{h}{n_2} \right) \right|^2.$$

By (1), the sum in question is

$$\ll N \sum_{n_1 \leq X} \frac{X}{n_1} Y \log^2 X \ll (NXY \log^2 X) \prod_{p|2N} \left(1 - \frac{1}{p} \right)^{-1}$$

which is the desired result.

LEMMA 2. Let d and a be fixed integers, with $a \equiv 1 \pmod{4}$, $(ad, 4N) = 1$. Then,

$$\left| \sum_{n \leq U, n_2 d \neq b^2} \frac{a(n)}{n} f_Y(nd; a) \right| \ll (Ud)^{1/2} Y^{1/2} \log Y \log(Ud).$$

Proof. By partial summation and Lemma 1,

$$\sum'_{m \leq U} \frac{1}{\sqrt{m}} \left| \sum_{\substack{h \leq Y \\ h \equiv a \pmod{4N}}} \left(\frac{h}{m} \right) \right|^2 \ll U^{1/2} Y \log^2 U.$$

Let $\chi_0^{(a)}$ denote the principal character mod a . The sum in question is

$$\begin{aligned} & \sum_{n \leq U, n_2 d \neq b^2} \frac{a(n)}{n} \sum_{\substack{0 < -D \leq Y \\ D \equiv a \pmod{4N}}} \left(\frac{D}{nd} \right) \sum_{j^2 | D} \mu(j) \\ &= \sum_{j^2 \leq Y} \mu(j) \sum_{n \leq U, n_2 d \neq b^2} \frac{a(n)}{n} \chi_0^{(j)}(nd) \sum_{\substack{0 < -h \leq Y/j^2 \\ h \equiv aj^2 \pmod{4N}}} \left(\frac{h}{nd} \right) \\ &\ll \sum_{j^2 \leq Y} \left(\sum_{n \leq U} \frac{|a(n)|^2}{n^{3/2}} \right)^{1/2} \left(\sum_{n \leq U, n_2 d \neq b^2} \frac{1}{\sqrt{n}} \left| \sum_{\substack{0 < -h \leq Y/j^2 \\ h \equiv aj^2 \pmod{4N}}} \left(\frac{h}{nd} \right) \right|^2 \right)^{1/2} \\ &\ll U^{1/4} d^{1/4} \sum_{j^2 \leq Y} \left(\sum'_{m \leq Ud} \frac{1}{\sqrt{m}} \left| \sum_{\substack{0 < -h \leq Y/j^2 \\ h \equiv aj^2 \pmod{4N}}} \left(\frac{h}{m} \right) \right|^2 \right)^{1/2} \\ &\ll U^{1/2} d^{1/2} Y^{1/2} (\log Y) (\log Ud), \end{aligned}$$

as desired.

LEMMA 3. For $\Re s = -\eta$, $0 < \eta < 1/2$, $a \equiv 1 \pmod{4}$, $(ad, 4N) = 1$,

$$\sum_{n \leq U, n_2 d \neq b^2} \frac{a(n) \chi_0^{(j)}(nd)}{n^{1-s}} \sum_{\substack{0 < -h \leq Y \\ h \equiv a \pmod{4N}}} \left(\frac{h}{nd} \right) \ll U^{1/2-\eta} d^{1/2} Y^{1/2} \log Ud.$$

Proof. We apply Cauchy's inequality: A

$$\begin{aligned} & \left(\sum_{n \leq U} \frac{|a(n)|^2}{n^{3/2+\eta}} \right)^{1/2} \left(\sum_{n \leq U, n_2 d \neq b^2} \frac{1}{n^{1/2+\eta}} \left| \sum_{\substack{0 < -h \leq Y \\ h \equiv a \pmod{4N}}} \left(\frac{h}{nd} \right) \right|^2 \right)^{1/2} \\ & \ll d^{1/2} U^{1/2-\eta} Y^{1/2} \log Ud \end{aligned}$$

by partial summation.

LEMMA 4. Under the same conditions as in Lemma 3,

$$\begin{aligned} & \sum_{n \leq U, n_2 d \neq b^2} \frac{a(n)}{n^{1-s}} \chi_0^{(j)}(nd) \sum_{\substack{0 < -h \leq Y \\ h \equiv a \pmod{4N}}} \left(\frac{h}{nd} \right) h^{-2s} \\ & \ll (|s| + 1) d^{1/2} U^{1/2-\eta} Y^{1/2+2\eta} \log Ud. \end{aligned}$$

Proof. Apply partial summation to Lemma 3.

LEMMA 5. If $\Re s = -\eta$, $0 < \eta < 1/2$, $a \equiv 1 \pmod{4}$, $(ad, 4N) = 1$, then

$$\sum_{n \leq U, n_2 d \neq b^2} \frac{a(n)}{n^{1-s}} f_Y(nd, -2s; a) \ll (|s| + 1) d^{1/2} U^{1/2-\eta} Y^{1/2+2\eta} \log Y \log Ud.$$

Proof. The sum in question is

$$\begin{aligned} & \sum_{n \leq U, n_2 d \neq b^2} \frac{a(n)}{n^{1-s}} \sum_{\substack{0 < -D \leq Y \\ D \equiv a \pmod{4N}}} \left(\frac{D}{nd} \right) \sum_{j^2 | D} \mu(j) D^{-2s} \\ & = \sum_{j^2 \leq Y} \mu(j) j^{-4s} \sum_{n \leq U, n_2 d \neq b^2} \frac{a(n) \chi_0^{(j)}(nd)}{n^{1-s}} \sum_{\substack{0 < -h \leq Y/j^2 \\ h \equiv aj^2 \pmod{4N}}} \left(\frac{h}{nd} \right) h^{-2s} \\ & \ll \sum_{j^2 \leq Y} j^{4\eta} (|s| + 1) (d^{1/2} U^{1/2-\eta} \log Ud) \frac{Y^{1/2+2\eta}}{j^{1+4\eta}} \end{aligned}$$

by Lemma 4. Summing over j gives the desired result.

Putting the above lemmas together proves:

LEMMA 6. For $\Re s = -\eta$, $0 < \eta < 1/2$, $a \equiv 1 \pmod{4}$, $(ad, 4N) = 1$,

(i)

$$\int_{(-\eta)} \sum_{n \leq U, n_2 d \neq b^2} \frac{a(n)}{n^{1-s}} \tilde{f}_Y(nd, -2s; a) \zeta(1+2s) \frac{\Gamma(1-s)}{\Gamma(1+s)} x^s \Gamma(s) ds \\ \ll x^{-\eta} d^{1/2} U^{1/2-\eta} Y^{1/2+2\eta} \log Ud,$$

and

(ii)

$$\int_{(-\eta)} \sum_{n \leq U, n_2 d \neq b^2} \frac{a(n)}{n^{1-s}} f_Y(nd, -2s; a) \zeta(1+2s) \frac{\Gamma(1-s)}{\Gamma(1+s)} x^s \Gamma(s) ds \\ \ll x^{-\eta} d^{1/2} U^{1/2-\eta} Y^{1/2+2\eta} \log Y \log Ud.$$

We now proceed to handle the terms corresponding to $n > U$.

LEMMA 7. For $\Re s = -\eta$, $\eta > 1/2$, $(ad, 4N) = 1$,

$$\sum_{n > U, n_2 d \neq b^2} \frac{a(n)}{n^{1-s}} \chi_0^{(j)}(nd) \sum_{\substack{0 < -h \leq Y \\ h \equiv a \pmod{4N}}} \left(\frac{h}{nd} \right) \ll U^{1/2-\eta} d^{1/2} Y^{1/2} \log Ud.$$

Proof. By Cauchy's inequality, the sum is bounded by

$$\left(\sum_{n > U} \frac{|a(n)|^2}{n^{3/2+\eta}} \right)^{1/2} \left(\sum_{n > U, n_2 d \neq b^2} \frac{1}{n^{1/2+\eta}} \left| \sum_{\substack{0 < -h \leq Y \\ h \equiv a \pmod{4N}}} \left(\frac{h}{nd} \right) \right|^2 \right)^{1/2} \\ \ll U^{1/4-\eta/2} d^{1/4+\eta/2} \left(\sum'_{m > Ud} \frac{1}{m^{1/2+\eta}} \left| \sum_{\substack{0 < -h \leq Y \\ h \equiv a \pmod{4N}}} \left(\frac{h}{m} \right) \right|^2 \right)^{1/2} \\ \ll U^{1/4-\eta/2} d^{1/4+\eta/2} \left(\int_{Ud}^{\infty} \frac{Y \log^2 t}{t^{1/2+\eta}} dt \right)^{1/2} \\ \ll U^{1/2-\eta} d^{1/2} Y^{1/2} \log Ud,$$

as desired.

LEMMA 8. Under the conditions of Lemma 7,

$$\sum_{n>U, n_2d \neq b^2} \frac{a(n)}{n^{1-s}} \chi_0^{(j)}(nd) \sum_{\substack{0 < -h \leq Y \\ h \equiv a \pmod{4N}}} \left(\frac{h}{nd} \right) h^{-2s} \\ \ll (|s| + 1) Y^{1/2+2\eta} d^{1/2} U^{1/2-\eta} \log Ud.$$

Proof. Apply partial summation to Lemma 7.

LEMMA 9. If $\Re s = -\eta$, $\eta > 1/2$, $a \equiv 1 \pmod{4}$, $(ad, 4N) = 1$, then

$$\sum_{n>U, n_2d \neq b^2} \frac{a(n)}{n^{1-s}} f_Y(nd, -2s; a) \ll (|s| + 1) d^{1/2} U^{1/2-\eta} Y^{1/2+2\eta} \log Y \log Ud.$$

Proof. The proof is analogous to that of Lemma 5, except we use Lemma 8 instead of Lemma 4.

Therefore, we deduce by putting these lemmas together:

LEMMA 10. For $\Re s = -\eta$, $1 > \eta > 1/2$, $a \equiv 1 \pmod{4}$, $(ad, 4N) = 1$,

(i)

$$\int_{(-\eta)} \sum_{n>U, n_2d \neq b^2} \frac{a(n)}{n^{1-s}} \tilde{f}_Y(nd, -2s; a) \zeta(1+2s) \frac{\Gamma(1-s)}{\Gamma(1+s)} x^s \Gamma(s) ds \\ \ll x^{-\eta} d^{1/2} U^{1/2-\eta} Y^{1/2+2\eta} \log Ud,$$

and

(ii)

$$\int_{(-\eta)} \sum_{n>U, n_2d \neq b^2} \frac{a(n)}{n^{1-s}} f_Y(nd, -2s; a) \zeta(1+2s) \frac{\Gamma(1-s)}{\Gamma(1+s)} x^s \Gamma(s) ds \\ \ll x^{-\eta} d^{1/2} U^{1/2-\eta} Y^{1/2+2\eta} \log Y \log Ud.$$

This is the series of lemmas needed. The next section establishes a lemma which refines the above estimates.

3. Further lemmas

We begin by proving:

LEMMA 11.

$$\sum_e \frac{1}{e} \exp(-ne^2/X) = \begin{cases} \frac{1}{2} \log \frac{X}{n} + \frac{1}{2} \gamma + \mathbf{O}((n/X)^{1+\varepsilon}) & \text{if } n < \frac{1}{2} X \\ \mathbf{O}(\exp(-n/X)) & \text{otherwise.} \end{cases}$$

Proof. This follows easily by partial summation (see, for example, Jutila [3, Lemma 1]).

LEMMA 12. *There is a $\rho > 0$ so that if $X \leq Y(\log Y)^{1+\rho}$, $a \equiv 1 \pmod{4}$, $(a, 4N) = 1$, then,*

$$\sum_{\substack{m, e \\ (m, j) = 1}} \frac{a(m)}{me} \tilde{f}_Y(m; a) \exp(-me^2/X) \ll Y \log^2 Y,$$

for any $j \geq 1$.

Remark. Note that in this sum we are not restricting to fundamental discriminants.

Proof. We split the sum into two parts corresponding to m_2 a square and m_2 not a square.

For the first part, it is

$$\begin{aligned} &= \sum_e \frac{1}{e} \sum_{(m_1 m_2, j) = 1} \frac{a(m_1 m_2^2)}{m_1 m_2^2} \exp(-m_1 m_2^2 e^2 / X) \\ &\quad \times \left\{ \left(\frac{a}{m_1} \right) \frac{Y}{4N} \frac{\phi(m_2)}{m_2} + O(d(m_2)) \right\}. \end{aligned}$$

Using Lemma 11, we see that the above sum is

$$\ll Y \sum_{m < \frac{1}{2}X} \frac{|a(m_1 m_2^2)|}{m_1 m_2^2} \log \frac{X}{m} + Y \sum_{m > \frac{1}{2}X} \frac{|a(m_1 m_2^2)|}{m_1 m_2^2} d(m) \exp(-m/X)$$

which is

$$\ll Y(\log X)^2.$$

For the second part, we use the Pólya-Vinogradov inequality to see that it is

$$\ll \sum_m \frac{|a(m)|}{m} m^{1/2} \log m \sum_e \frac{1}{e} \exp(-me^2/X).$$

Using Lemma 11 for the inner sum, we see that it is

$$\ll \sum_{m < \frac{1}{2}X} \frac{|a(m)|}{m} m^{1/2} \log m \left\{ \log \frac{X}{m} \right\} + \sum_{m > \frac{1}{2}X} \frac{|a(m)|}{m} m^{1/2} \log m \exp(-m/X)$$

and by partial summation and Lemma 17, this is

$$\ll X(\log X)^{1-\rho},$$

for some $\rho > 0$. This completes the proof.

LEMMA 13. Let $a \equiv 1 \pmod{4}$ with $(a, 4N) = 1$ and a equal to a square mod $4N$. Then,

$$\sum_{\substack{0 < -D \leq Y \\ D \equiv a \pmod{4N}}} L'_D(1) \ll Y \log^2 Y.$$

Proof. Consider

$$\frac{1}{2\pi i} \int_{(\sigma)} L_D(1+s) \zeta(1+2s) X^s \Gamma(s) ds, \quad \sigma > 1/2.$$

This is

$$\sum_{n,m} \frac{a(n)}{n} \left(\frac{D}{n} \right) \frac{1}{m} \exp(-nm^2/X).$$

(Note that this series converges absolutely.) By the Phragmén-Lindelöf theorem [6],

$$|L_D(1+s)| \ll (|t|+2)^{1/2-\sigma} (\log(|t|+2))^2$$

uniformly for $-1/2 - \varepsilon \leq \sigma \leq 1/2 + \varepsilon$. (The implied constant depends on D .) We can therefore move the line of integration to $\Re s = -\eta \geq -1/2 - \varepsilon$.

Moving the line of integration to $\Re s = -\eta$, we obtain

$$\begin{aligned} \sum_{n,m} \frac{a(n)}{n} \left(\frac{D}{n} \right) \frac{1}{m} \exp(-nm^2/X) \\ = \frac{1}{2} L'_D(1) + \frac{1}{2\pi i} \int_{(-\eta)} L_D(1+s) \zeta(1+2s) X^s \Gamma(s) ds. \end{aligned}$$

Writing

$$L_D(1+s) = L_{D_0}(1+s) \sum_{d|\delta^2} \frac{\tilde{\mu}(d)}{d^{1+s}} \left(\frac{D_0}{d} \right)$$

where D_0 is a fundamental discriminant, we can rewrite the integral as

$$\frac{1}{2\pi i} \sum_{d|\delta^2} \frac{\tilde{\mu}(d)}{d} \int_{(-\eta)} L_{D_0}(1+s) \left(\frac{D_0}{d} \right) \zeta(1+2s) (X/d)^s \Gamma(s) ds.$$

Applying the functional equation to $L_{D_0}(1+s)$, we find that the above is equal to

$$\begin{aligned} \frac{w \chi_{D_0}(-N)}{2\pi i} \sum_{d|\delta^2} \frac{\tilde{\mu}(d)}{d} \int_{(-\eta)} \sum_{n=1}^{\infty} \frac{a(n)}{n^{1-s}} \left(\frac{D_0}{nd} \right) \zeta(1+2s) |D_0|^{-2s} A^{-2s} \\ \times \frac{\Gamma(1-s)}{\Gamma(1+s)} (X/d)^s \Gamma(s) ds. \end{aligned}$$

We sum this over $\delta^2 \leq Y$, $(\delta, 4N) = 1$, $0 < -D_0 \leq Y/\delta^2$, $D_0\delta^2 \equiv a \pmod{4N}$. Recall that we have set

$$\tilde{f}_Y(n, s; a) = \sum_{\substack{0 < -D \leq Y \\ D \equiv a \pmod{4N}}} \left(\frac{D}{n} \right) |D|^s, \quad D \text{ unrestricted},$$

$$f_Y(n, s; a) = \sum_{\substack{0 < -D_0 \leq Y \\ D_0 \equiv a \pmod{4N}}} \left(\frac{D_0}{n} \right) |D_0|^s, \quad D_0 \text{ fundamental},$$

$$\tilde{f}_Y(n; a) = \tilde{f}_Y(n, 0; a), \quad f_Y(n; a) = f_Y(n, 0; a).$$

Then,

$$\begin{aligned} (\star) \quad & \sum_{n, m} \frac{a(n)}{n} \tilde{f}_Y(n; a) \frac{1}{m} \exp(-nm^2/X) \\ &= \frac{1}{2} \sum_{\substack{0 < -D \leq Y \\ D \equiv a \pmod{4N}}} L'_D(1) - \frac{1}{2\pi i} \sum_{\substack{\delta^2 \leq Y \\ (\delta, 4N)=1}} \sum_{d|\delta^2} \frac{\bar{\mu}(d)}{d} \int_{(-\eta)} \sum_{n=1}^{\infty} \frac{a(n)}{n^{1-s}} \\ & \quad \times f_{Y/\delta^2}(nd, -2s; a\bar{\delta}^2) \zeta(1+2s) \\ & \quad \times \frac{\Gamma(1-s)}{\Gamma(1+s)} A^{-2s} (X/d)^s \Gamma(s) ds. \end{aligned}$$

Here $\bar{\delta}\delta \equiv 1 \pmod{4N}$. The sum on the left side of (\star) is seen to be

$$\ll Y(\log Y)^2$$

by Lemma 12, provided we take $X \leq Y(\log Y)^{1+\rho}$. Now we deal with the integral on the right side of (\star) . The integral is first broken up according to whether n_2d is a square or not. If n_2d is not a square, then splitting the series at $n \leq Y$ and moving the integral to $\Re s = -\eta_1$, $0 < \eta_1 < 1/2$, we can utilise Lemma 6(ii) to get an estimate

$$\sum_{\delta^2 \leq Y} \sum_{d|\delta^2} \frac{|\bar{\mu}(d)|}{d} d^{1/2} \frac{Y}{\delta^{1+3\eta_1}} \left(\frac{Y}{X} \right)^{\eta_1} \log^2 Y \ll Y(Y/X)^{\eta_1} \log^2 Y.$$

The series corresponding to $n \geq Y$ is similarly estimated by moving to $\Re s = -\eta_2$, $\eta_2 > 1/2$ and using Lemma 10(ii). The final contribution when n_2d is not a square is therefore

$$Y \left\{ \left(\frac{Y}{X} \right)^{\eta_1} + \left(\frac{Y}{X} \right)^{\eta_2} \right\} \log^2 Y.$$

This term is $\ll Y \log^2 Y$ if $Y \leq X$. Now we consider the contribution when $n_2 d$ is a square.

The series in the integral is seen to be

$$\sum_{n_1, n_2} \frac{a(n_1 n_2)}{(n_1 n_2)^{1-s}} \left(\frac{a}{n_1} \right) f_{Y/\delta^2}(n_2 d, -2s, a\bar{\delta}^2).$$

Let

$$H_d(s) = \sum_{\substack{n_1, n_2 \\ n_2 d = b^2}} \frac{a(n_1 n_2)}{(n_1 n_2)^s} \left(\frac{a}{n_1} \right) \prod_{p|4Nn_2 d} \left(1 + \frac{1}{p} \right)^{-1}.$$

We see that this is equal to

$$\begin{aligned} & \sum_{\substack{n_1, n_2 \\ n_2 d = b^2}} \frac{a(n_1) a(n_2)}{(n_1 n_2)^s} \left(\frac{a}{n_1} \right) \prod_{p|4Nn_2 d} \left(1 + \frac{1}{p} \right)^{-1} \\ &= \left(\sum_{p|n \Rightarrow p|2N} \frac{a(n_1)}{n_1^s} \left(\frac{a}{n_1} \right) \right) \left(\sum_{n_2 d = b^2} \frac{a(n_2)}{n_2^s} \prod_{p|4Nn_2 d} \left(1 + \frac{1}{p} \right)^{-1} \right). \end{aligned}$$

Thus, in order to estimate the integral, we first need an estimate for $H_d(s)$. Since the first sum is uniformly bounded in the region under consideration, we need only consider the second Dirichlet series. If $d = d_0 d_1^2$, d_0 squarefree, then

$$\sum_{n_2 d = b^2} \frac{a(n_2)}{n_2^s} \prod_{p|4Nn_2 d} \left(1 + \frac{1}{p} \right)^{-1} = \sum_m \frac{a(d_0 m^2)}{d_0^s m^{2s}} \prod_{p|4Nmd} \left(1 + \frac{1}{p} \right)^{-1}$$

where the sum over m is such that $(m, 2N) = 1$. Let us now estimate the integral. For this purpose, let us define

$$F_d(s) = \sum_{n=1}^{\infty} \frac{a(d_0 n^2)}{(d_0 n^2)^s} \prod_{p|4Nnd} \left(1 + \frac{1}{p} \right)^{-1}.$$

By factoring $F_d(s)$ as an Euler product and using simple estimates, we find that

$$|F_d(s)| \ll c^{\nu(d)} d_0^{1/2-\sigma} |L(2s, \text{Sym}^2)\zeta(4s-2)^{-1}|,$$

for $\sigma > 3/4$. Here, c is an absolute positive constant (we can take $c = 200$ for example) and $\nu(d)$ is the number of prime factors of d .

We are now ready to estimate the integral when $n_2 d$ is a square. We move this integral to $\eta = 1/\log Y$ and evaluate it using the above estimate for $F_d(s)$.

By an easy variant of Jutila's result [3, Lemma 1], we find the integral is

$$\begin{aligned} & \frac{1}{2\pi i} \int_{(-\eta)} \sum_{\substack{n=1 \\ n_2 d = b^2}}^{\infty} \frac{a(n)}{n^{1-s}} \left(\frac{a}{n_1} \right) A^{-2s} \frac{\Gamma(1-s)}{\Gamma(1+s)} \zeta(1+2s) (X/d)^s \Gamma(s) \\ & \times \left\{ \frac{6}{\pi^2} \frac{1}{\phi(4N)} \prod_{p|4Nn_2d} \left(1 + \frac{1}{p} \right)^{-1} \frac{(Y/\delta^2)^{1-2s}}{1-2s} \right. \\ & \quad \left. + O\left((|s|+1) d(4Nn_2d) \frac{Y^{1/2+2\eta}}{\delta^{1+4\eta}} \right) \right\} ds. \end{aligned}$$

The error term presents no problem. The main term is

$$\ll \frac{Y}{\delta^2} (d/X)^{\eta} c^{\nu(d)} (\log Y)^2.$$

Summing over $d|\delta^2$ and $\delta^2 \leq Y$, we obtain a total estimate of

$$\ll Y \sum_{\delta^2 \leq Y} \frac{1}{\delta^2} \sum_{d|\delta^2} \frac{|\tilde{\mu}(d)|}{d} (d/X)^{\eta} c^{\nu(d)} (\log Y)^2 \ll Y (\log Y)^2.$$

This completes the proof of Lemma 13.

Utilising Lemma 13, we can now derive an alternate estimate for the sum

$$\sum_{m,e} \frac{a(m)}{me} \tilde{f}_Y(m; a) \exp\left(-\frac{me^2}{X}\right).$$

LEMMA 14. If $Y \leq X$, $a \equiv 1 \pmod{4}$, $(ad, 4N) = 1$, and any $j \geq 1$,

$$(i) \quad \sum_{\substack{m,e \\ (m,j)=1}} \frac{a(m)}{me} \tilde{f}_Y(md; a) \exp(-me^2/X) \ll d^{1/2} Y \log^2 Y.$$

Also

$$(ii) \quad \sum'_{\substack{m,e \\ (m,j)=1}} \frac{a(m)}{me} \tilde{f}_Y(md; a) \exp(-me^2/X) \ll d^{1/2} Y (\log^2 Y + \log X),$$

where the prime on the sum indicates that $m_2 d$ is not a square.

Proof. Consider equation (\star) in the proof of Lemma 13, which shows that under the hypothesis $Y \leq X$, the integral in (\star) is $O(Y \log^2 Y)$. Lemma 13 itself asserts that the same is true of the sum of the $L'_D(1)$. Our assertion follows for

$j = 1$ and $d = 1$. In general, we consider an analogue of Lemma 13 in which we multiply both sides of the original equation by (D/d) and only sum over $D = \delta^2 D_0$ with $j \mid \delta$. The estimates proceed exactly as before and we obtain assertion (i) of the lemma. For (ii), we need only observe that the contribution from those m, e with $m_2 d$ a square is $\ll d^{1/2} Y(\log X)$.

LEMMA 15. If $a \equiv 1 \pmod{4}$, $(ad, 4N) = 1$, then

$$\sum'_{\substack{m, e \\ (m, j)=1}} \frac{a(m)}{me} \tilde{f}_Y(md; a) \exp(-me^2/X) \ll d^{1/2} X^{1/2} Y^{1/2} \log Xd$$

where the prime indicates that the sum ranges over values of m such that $m_2 d$ is not a square.

Proof. We have to estimate

$$\sum'_{\substack{m, e \\ (m, j)=1}} \frac{a(m)}{me} \exp(-me^2/X) \sum_{\substack{0 < -h \leq Y \\ h \equiv a \pmod{4N}}} \left(\frac{h}{md} \right).$$

We bring the summation over e inside and use Lemma 11. If we truncate the sum over m at $\frac{1}{2}X$, then it is

$$\ll \sum_{\substack{m < \frac{1}{2}X \\ m_2 d \neq b^2}} \frac{|a(m)|}{m} \left| \sum_{\substack{0 < -h \leq Y \\ h \equiv a \pmod{4N}}} \left(\frac{h}{md} \right) \right| \left(\log \frac{X}{m} + O(1) \right),$$

and by Lemma 1 and the method of Lemma 2, this is

$$\ll (Xd)^{1/4} \left\{ (Xd)^{1/2} Y (\log Xd)^2 \right\}^{1/2}.$$

The sum over $m > \frac{1}{2}X$ is estimated in the same way. It is

$$\ll \sum_{\substack{m > \frac{1}{2}X \\ m_2 d \neq b^2}} \frac{|a(m)|}{m} \exp(-m/X) \left| \sum_{\substack{0 < -h \leq Y \\ h \equiv a \pmod{4N}}} \left(\frac{h}{md} \right) \right|$$

which is

$$\ll (Xd)^{1/4} \left\{ (Xd)^{1/2} Y (\log Xd)^2 \right\}^{1/2}.$$

In both cases, the estimate simplifies to

$$\ll (dXY)^{1/2} (\log Xd).$$

This proves the lemma.

We shall use this to prove the following crucial and penultimate lemma.

LEMMA 16. Take $a \equiv 1 \pmod{4}$, $(ad, 4N) = 1$. For all X satisfying $X \geq Y \log^{-B} Y$,

$$\sum'_{m,e} \frac{a(m)}{me} f_Y(md; a) \exp(-me^2/X) \ll_B d^{1/2} X^{1/2} Y^{1/2} (\log X) (\log \log Y)$$

where the prime indicates that the sum ranges over values of m such that $m_2 d$ is not a square.

Proof. We have

$$\begin{aligned} & \sum'_{m,e} \frac{a(m)}{me} f_Y(md; a) \exp(-me^2/X) \\ &= \sum_{\substack{j^2 \leq Y \\ (j,d)=1}} \mu(j) \sum'_{\substack{m,e \\ (m,j)=1}} \frac{a(m)}{me} \exp(-me^2/X) \sum_{\substack{0 < -h \leq Y/j^2 \\ h \equiv aj^2 \pmod{4N}}} \left(\frac{h}{md} \right). \end{aligned}$$

The inner sum is what we have denoted by $f_{Y/j^2}(md; aj^2)$. The sum of the terms with $j < \log^B Y$ can be estimated by Lemma 15 and it is seen to be

$$\ll \sum_{j < \log^B Y} \frac{(dXY)^{1/2}}{j} \log Xd$$

which is

$$d^{1/2} X^{1/2} Y^{1/2} \log X \log \log Y.$$

For $j > \log^B Y$, we use the second part of Lemma 14 to get

$$\sum_{j > \log^B Y} d^{1/2} \frac{Y}{j^2} (\log^2 Y + \log X) \ll d^{1/2} Y \log^{-B+2} Y.$$

Since $X \geq Y \log^{-B} Y$, this proves the lemma.

The final lemma is an estimate for a weighted average of the $a(n)$ which will be needed in some of the error term estimates.

LEMMA 17. There is a $\rho > 0$ so that

$$\sum_{n \leq x} \frac{|a(n)|}{\sqrt{n}} \ll x (\log x)^{-\rho}$$

and

$$\sum_{n \leq x} \frac{|a(n)|}{\sqrt{n}} d(n) \ll x (\log x)^{1-2\rho}.$$

Proof. The first estimate is due to Rankin [10]. (It should be remarked that Rankin proved the estimate for $N = 1$. However, his proof carries over for the general level in view of Shahidi's result [12].)

For the second estimate, let $b_n = |a(n)|/n^{1/2}$ and consider the Dirichlet series $F(s) = \sum_{n=1}^{\infty} b_n \mathbf{d}(n)/n^s$. Set $Q(s) = \sum_{n=1}^{\infty} \mu^2(n) b_n \mathbf{d}(n)/n^s$. We are interested in bounding the partial sums of $F(s)$. Consider first the partial sums of $Q(s)$, namely,

$$\sum_{n \leq x} \mu^2(n) b_n \mathbf{d}(n).$$

The coefficients of $Q(s)$ are dominated by the coefficients of $(\sum_{n=1}^{\infty} b_n/n^s)^2$. By the first part of the lemma,

$$\sum_{n \leq x} b_n \ll x/(\log x)^{\rho}.$$

Hence,

$$\sum_{mn \leq x} b_m b_n \ll \sum_{n \leq \sqrt{x}} b_n \sum_{m \leq x/n} b_m \ll \sum_{n \leq \sqrt{x}} b_n (x/n) (\log x)^{-\rho} \ll x (\log x)^{1-2\rho}$$

by partial summation.

Now set $F(s) = Q(s)R(s)$. Write $R(s) = \sum c(n)n^{-s}$. We see that as the Euler product of $R(s)$ converges absolutely for $\operatorname{Re}(s) > 1/2$, the Dirichlet series $\sum c(n)/n$ is absolutely convergent. Using these facts, we see that

$$\begin{aligned} \sum_{n \leq x} b_n \mathbf{d}(n) &= \sum_{me \leq x} c(m) \mu^2(e) b_e \mathbf{d}(e) \\ &\ll \sum_{m \leq x} c(m) \frac{2x}{m} \left(\log \frac{2x}{m} \right)^{1-2\rho} \\ &\ll x (\log x)^{1-2\rho}. \end{aligned}$$

This proves the lemma.

4. The main theorems

Now consider

$$\frac{1}{2\pi i} \int_{(\sigma)} L_{D_0}(1+s) \zeta(1+2s) \sum_{d|\delta^2} \frac{\tilde{\mu}(d)}{d^{1-s}} \left(\frac{D_0}{d} \right)^{X^s} \Gamma(s) \delta^{-4s} ds.$$

This is

$$\sum_{d|\delta^2} \frac{\tilde{\mu}(d)}{d} \sum_{m,e} \frac{a(m)}{me} \left(\frac{D_0}{md} \right) \exp(-me^2\delta^4/Xd).$$

On the other hand, moving the line of integration to the left and picking up the residue at $s = 0$, we obtain

$$\frac{1}{2}L'_D(1) + \frac{1}{2\pi i} \int_{(-\eta)} L_{D_0}(1+s)\zeta(1+2s) \sum_{d|\delta^2} \frac{\tilde{\mu}(d)}{d^{1-s}} \left(\frac{D_0}{d} \right) X^s \Gamma(s) \delta^{-4s} ds.$$

Writing the functional equation for $L_{D_0}(s)$, we see that the above integral is

$$\frac{w\chi_{D_0}(-N)}{2\pi i} \int_{(-\eta)} |D|^{-2s} L_D(1-s)\zeta(1+2s) \frac{\Gamma(1-s)}{\Gamma(1+s)} A^{-2s} X^s \Gamma(s) ds.$$

Let Y be such that $Y \log^{-B} Y \leq X \leq Y(\log Y)^{1+\nu}$ where $0 < \nu < \rho$, with ρ as in Lemma 17. Now we sum the entire expression above over $\delta^2 \leq Y$, $0 < -D_0 \leq Y/\delta^2$, $D_0\delta^2 \equiv 1 \pmod{4N}$ and obtain the expression:

$$\begin{aligned} & \frac{1}{2} \sum_{\substack{0 < -D \leq Y \\ D \equiv 1 \pmod{4N}}} L'_D(1) \\ &= \sum_{\substack{\delta^2 \leq Y \\ (\delta, 4N)=1}} \sum_{d|\delta^2} \frac{\tilde{\mu}(d)}{d} \sum_{n=1}^{\infty} \sum_{me^2=n} \frac{a(m)}{me} f_{Y/\delta^2}(md; \bar{\delta}^2) \exp(-n\delta^4/Xd) \\ & \quad + \frac{1}{2\pi i} \int_{(-\eta)} \sum_{\substack{m=1 \\ m=m_1m_2}}^{\infty} \frac{a(m)}{m^{1-s}} \tilde{f}_Y(m, -2s; 1) \zeta(1+2s) \\ & \quad \times \frac{\Gamma(1-s)}{\Gamma(1+s)} A^{-2s} X^s \Gamma(s) ds, \end{aligned}$$

for $\eta > 1/2$.

The integral is split into two parts. In the first, the sum over m_2 is taken only over non-squares and, in the second, it ranges over squares.

Let us consider the first integral. It is easily estimated in the following way. Truncating the sum over m at U , we move the integral involving the part $m \leq U$ to $\Re s = -\eta_1$ where $\eta_1 < 1$. The part corresponding to $m > U$ is moved to $\Re s = -\eta_2$ where $\eta_2 > 1$. Choosing $U = Y^2/X$ and using the Pólya-Vinogradov inequality and Lemma 17, we see that the integral is $\ll X^{-1}Y^2(\log Y)^{1-\nu}$.

The contribution from the squares is easily handled. With a slight change in notation, it is

$$\frac{1}{2\pi i} \int_{(-\eta)} \sum_{m=1}^{\infty} \frac{a(m_1 m_2^2)}{(m_1 m_2^2)^{(1-s)}} \zeta(1+2s) \sum_D |D|^{-2s} \frac{\Gamma(1-s)}{\Gamma(1+s)} X^s A^{-2s} \Gamma(s) ds$$

where the sum over D is over

$$0 < -D \leq Y, \quad D \equiv 1 \pmod{4N}, \quad (D, m_2) = 1.$$

First, we move the integral to a line $-\eta$, with $0 < \eta < \frac{1}{2}$. By partial summation, it is easily seen that the sum over D is asymptotic to

$$\frac{1}{\phi(4N)} \frac{\phi(4Nm_2)}{4Nm_2} \frac{Y^{1-2s}}{1-2s} + O(Y^{2\eta} \mathbf{d}(m_2)(|s|+1)),$$

where, as before, $\mathbf{d}(m_2)$ is the number of divisors of m_2 . Inserting this into the integral, we find that the main term is

$$\frac{Y}{\phi(4N)} \frac{1}{2\pi i} \int_{(-\eta)} T(1-s) \left(\frac{X}{A^2 Y^2} \right)^s \frac{\Gamma(1-s)}{\Gamma(1+s)} \frac{1}{1-2s} \zeta(1+2s) \Gamma(s) ds$$

where

$$T(s) = \sum_{n_1, n_2} \frac{\phi(4Nn_2) a(n_1 n_2^2)}{4Nn_2 (n_1 n_2^2)^s}.$$

The error term is easily seen to be $O(Y^{2\eta} X^{-\eta})$. Now moving this integral to the right of $\Re s = 0$, and using the expansions

$$\frac{\Gamma(1-s)}{\Gamma(1+s)} \frac{1}{1-2s} = 1 + (2\gamma+2)s + (2\gamma^2+4\gamma+4)s^2 + \cdots,$$

$$\left(\frac{X}{A^2 Y^2} \right)^s = 1 + \left(\log \frac{X}{A^2 Y^2} \right) s + \frac{1}{2} \left(\log^2 \frac{X}{A^2 Y^2} \right) s^2 + \cdots$$

and

$$\zeta(1+2s) \Gamma(s) = \left\{ \frac{1}{2s} + \gamma + \cdots \right\} \left\{ \frac{1}{s} - \gamma + \cdots \right\} = \frac{1}{2s^2} + \frac{\gamma}{2s} + \cdots,$$

we find that it is

$$-\frac{Y}{\phi(4N)} \sum_{n_1, n_2} \frac{a(n_1 n_2^2)}{n_1 n_2^2} \frac{\phi(4Nn_2)}{4Nn_2} \left(\frac{1}{2} \log \frac{X}{Y^2} \right) + O(Y).$$

We see that this is

$$-\frac{1}{4} CY \log \frac{X}{Y^2} + O(Y).$$

Summarizing, we have proved that the integral is

$$-\frac{1}{4}CY \log \frac{X}{Y^2} + O(Y) + O\left(\frac{1}{X}Y^2(\log Y)^{1-\nu}\right).$$

Next, we consider the sum which we split into the contribution when m_2d is a perfect square and when m_2d is not a perfect square.

In the latter case, we estimate the sum as follows. First consider the contribution of terms in which $\delta > \log^A Y$, for some large $A > 0$. By using Lemma 2 and partial summation, we see that

$$\begin{aligned} & \sum_e \frac{1}{e} \sum'_m \frac{a(m)}{m} f_{Y/\delta^2}(md; \bar{\delta}^2) \exp(-me^2\delta^4/Xd) \\ & \ll \left(\frac{Xd}{\delta^4}d\right)^{1/2} \frac{Y^{1/2}}{\delta} (\log Y)(\log X). \end{aligned}$$

Thus the contribution of these terms to the sum is

$$\ll \sum_{\delta > \log^A Y} \sum_{d|\delta^2} \frac{|\tilde{\mu}(d)|}{d} \left(\frac{Xd}{\delta^4}d\right)^{1/2} \frac{Y^{1/2}}{\delta} (\log Y)(\log X)$$

and this is clearly $O(Y)$. For $\delta < \log^A Y$, we argue as above, except that we use the estimate of Lemma 16 in place of Lemma 2. We find that these terms contribute an amount

$$\ll X^{1/2}Y^{1/2} \log X \log \log X.$$

In the former case, it is

$$\sum_{\delta} \sum_{d|\delta^2} \frac{\tilde{\mu}(d)}{d} \sum_{n=1}^{\infty} \sum_{\substack{me^2=n \\ m_2d=r^2}} \frac{a(m)}{me} \left(\sum_{D_0} 1\right) \exp(-n\delta^4/Xd)$$

where the sum on δ ranges over

$$\delta^2 \leq Y, \quad (\delta, 4N) = 1$$

and the sum on D_0 ranges over

$$0 < -D_0 \leq \frac{Y}{\delta^2}, \quad D_0 \equiv \bar{\delta}^2 \pmod{4N}, \quad (D_0, m_2d) = 1,$$

$$D_0 \equiv 1 \pmod{4}, \quad D_0 \text{ fundamental.}$$

The sum over D_0 can be expressed as

$$\sum_{0 < -D_0 \leq Y/\delta^2} \mu^2(-D_0) \chi_0^{(m_2 d)}(D_0) \frac{1}{\phi(4N)} \sum_{\psi \bmod 4N} \bar{\psi}(\bar{\delta}^2) \psi(D_0)$$

which is equal to

$$\sum_{\psi \bmod 4N} \bar{\psi}(\bar{\delta}^2) \frac{1}{\phi(4N)} \sum_{0 < -D_0 \leq Y/\delta^2} \mu^2(-D_0) \chi_0^{(m_2 d)}(D_0) \psi(D_0).$$

Now we consider the inner sum,

$$\frac{1}{\phi(4N)} \sum_{0 < -D_0 \leq Y/\delta^2} \left(\sum_{a^2 | D_0} \mu(a) \right) \chi_0^{(m_2 d)}(D_0) \psi(D_0).$$

Rearranging, we find it is

$$\frac{1}{\phi(4N)} \sum_{a \leq \sqrt{Y}/\delta} \mu(a) \chi_0^{(m_2 d)}(a^2) \psi(a^2) \sum_{0 < -h \leq Y/\delta^2 a^2} \chi_0^{(m_2 d)}(h) \psi(h).$$

If ψ is nontrivial, then $\psi \chi_0^{(m_2 d)}$ is a nontrivial character of conductor dividing $4N$. Thus, by Pólya-Vinogradov, the innermost sum is $O(1)$. The whole sum is then

$$O\left(\frac{Y^{1/2}}{\delta} \frac{1}{\phi(4N)}\right).$$

On the other hand, if ψ is the trivial character mod $4N$, we get

$$\frac{1}{\phi(4N)} \sum_{a \leq \sqrt{Y}/\delta} \mu(a) \chi_0^{(m_2 d)}(a^2) \chi_0^{(4N)}(a^2) \left\{ \frac{Y}{\delta^2 a^2} \frac{\phi(4Nm_2 d)}{4Nm_2 d} + O(d(4Nm_2 d)) \right\}.$$

Inserting this information into our big sum, we find that it is

$$\begin{aligned} & \sum_{\substack{\delta^2 \leq Y \\ (\delta, 4N)=1}} \sum_{d|\delta^2} \frac{\tilde{\mu}(d)}{d} \sum_{n=1}^{\infty} \sum_{\substack{me^2=n \\ m_2 d=r^2}} \frac{a(m)}{me} \exp(-n\delta^4/Xd) \frac{1}{\phi(4N)} \\ & \times \left\{ \frac{Y}{\delta^2} \frac{6}{\pi^2} \prod_{p|4Nm_2 d} \left(1 - \frac{1}{p^2}\right)^{-1} \frac{\phi(4Nm_2 d)}{4Nm_2 d} \right. \\ & \left. + O\left(\frac{Y^{1/2}}{\delta} d(4Nm_2 d)\right) \right\}. \end{aligned}$$

We observe that the contribution from terms with $\delta > X^{1/5}$ is negligible.

Indeed, it is

$$\ll \sum_{\sqrt{Y} > \delta > X^{1/5}} \frac{Y^{1/2}}{\delta} \sum_{d|\delta^2} \frac{|\tilde{\mu}(d)|}{d} \sum_{m=1}^{\infty} \frac{|a(m)|}{m} \\ \times \left(\frac{Y^{1/2}}{\delta} + \mathbf{d}(4Nmd) \right) \left(\sum_e \frac{1}{e} \exp(-me^2\delta^4/Xd) \right).$$

We use Lemma 11 to estimate the sum over e .

The terms with $m > Xd/2\delta^4$ contribute an amount $\mathbf{O}(S)$ where

$$S = \sum_{\sqrt{Y} > \delta > X^{1/5}} \frac{Y^{1/2}}{\delta} \sum_{d|\delta^2} \frac{|\tilde{\mu}(d)|}{d} \sum_{m > Xd/2\delta^4} \frac{|a(m)|}{m} \\ \times \left(\frac{Y^{1/2}}{\delta} + \mathbf{d}(4Nmd) \right) \exp\left(-\frac{m\delta^4}{Xd}\right).$$

By Lemma 17 and partial summation, we have

$$\sum_{m > Xd/2\delta^4} \frac{|a(m)|}{m} \mathbf{d}(m) \exp\left(-\frac{m\delta^4}{Xd}\right) \ll \sqrt{\frac{Xd}{\delta^4}} \left(\log \frac{Xd}{\delta^4} \right).$$

Thus,

$$S \ll \sum_{\sqrt{Y} > \delta > X^{1/5}} \frac{Y^{1/2}}{\delta} \sum_{d|\delta^2} \left(\frac{Y^{1/2}}{\delta} + \mathbf{d}(4Nd) \right) \frac{|\tilde{\mu}(d)|}{d} \sqrt{\frac{Xd}{\delta^4}} \left(\log \frac{Xd}{\delta^4} \right)$$

and this is

$$\ll (XY)^{1/2} (\log X) \sum_{\delta > X^{1/5}} \frac{1}{\delta^3} \sum_{d|\delta^2} \left(\frac{Y^{1/2}}{\delta} + \mathbf{d}(4Nd) \right) \mathbf{d}(d).$$

Since $\mathbf{d}(n) \ll n^\varepsilon$, it follows that

$$S \ll YX^{-1/10+\varepsilon}.$$

Now consider the terms with $m < Xd/2\delta^4$. Again, by Lemma 11, the sum to be estimated is

$$\ll \sum_{\sqrt{Y} > \delta > X^{1/5}} \frac{Y^{1/2}}{\delta} \sum_{d|\delta^2} \frac{|\tilde{\mu}(d)|}{d} \sum_{m < Xd/2\delta^4} \left(\frac{Y^{1/2}}{\delta} + \mathbf{d}(4Nmd) \right) \frac{|a(m)|}{m} \\ \times \left(\log \frac{Xd}{m\delta^4} + \mathbf{O}(1) \right).$$

By an argument analogous to that used in the treatment of S we see that the above is also

$$\ll YX^{-1/10+\varepsilon}.$$

Thus, the sum to be considered is

$$\sum_{\substack{\delta \leq X^{1/5} \\ (\delta, 4N)=1}} \sum_{d|\delta^2} \frac{\tilde{\mu}(d)}{d} \sum_{n=1}^{\infty} \sum_{\substack{me^2=n \\ m_2d=r^2}} \frac{a(m)}{me} \exp(-n\delta^4/Xd) \frac{1}{\phi(4N)} \\ \times \left\{ \frac{Y}{\delta^2} \frac{6}{\pi^2} \prod_{p|4Nm_2d} \left(1 - \frac{1}{p^2}\right)^{-1} \frac{\phi(4Nm_2d)}{4Nm_2d} + O\left(\frac{Y^{1/2}}{\delta} d(4Nm_2d)\right) \right\}.$$

The error term is

$$\ll Y^{1/2} \sum_{\delta \leq X^{1/5}} \frac{1}{\delta} \sum_{d|\delta^2} \frac{d(d)^2}{\sqrt{d}} \left\{ \sum_{m \leq Xd/2\delta^4} \frac{d(m)|a(m)|}{m} \log \frac{Xd}{m\delta^4} \right. \\ \left. + O\left(\sum_{m \geq Xd/2\delta^4} \frac{d(m)|a(m)|}{m} \exp\left(-\frac{m\delta^4}{Xd}\right) \right) \right\}.$$

Using Lemma 17 and partial summation, we see that

$$\sum_{m \leq Xd/2\delta^4} \frac{d(m)|a(m)|}{m} \log \frac{Xd}{m\delta^4} \ll \sqrt{\frac{Xd}{\delta^4}} \log X$$

and

$$\sum_{m \geq Xd/2\delta^4} \frac{d(m)|a(m)|}{m} \exp\left(-\frac{m\delta^4}{Xd}\right) \ll \sqrt{\frac{Xd}{\delta^4}} \log X.$$

Therefore, the error term is

$$\ll Y^{1/2} \sum_{\delta \leq X^{1/5}} \frac{1}{\delta} \sum_{d|\delta^2} \frac{d(d)^2}{\sqrt{d}} \frac{X^{1/2}d^{1/2}}{\delta^2} \log X \ll X^{1/2}Y^{1/2} \log X.$$

Thus, the sum reduces to

$$\frac{6Y}{\pi^2\phi(4N)} \sum_{\substack{\delta \leq X^{1/5} \\ (\delta, 4N)=1}} \frac{1}{\delta^2} \sum_{d|\delta^2} \frac{\tilde{\mu}(d)}{d} \sum_{n=1}^{\infty} \sum_{\substack{me^2=n \\ m_2d=r^2}} \frac{a(m)}{me} \exp(-n\delta^4/Xd) \\ \times \prod_{p|4Nm_2d} \left(1 + \frac{1}{p}\right)^{-1} + O(X^{1/2}Y^{1/2} \log X).$$

Now the sum over n is analysed as follows:

$$\begin{aligned} \sum_{n=1}^{\infty} \sum_{\substack{me^2=n \\ m_2d=r^2}} \frac{a(m)}{me} \prod_{p|4Nm_2d} \left(1 + \frac{1}{p}\right)^{-1} \exp(-n/X) \\ = \frac{1}{2\pi i} \int_{(2)} G_d(1+s) \zeta(1+2s) X^s \Gamma(s) ds \end{aligned}$$

where

$$G_d(s) = \sum_{\substack{m=1 \\ m_2d=r^2}}^{\infty} \frac{a(m)}{m^s} \prod_{p|4Nm_2d} \left(1 + \frac{1}{p}\right)^{-1}.$$

By the argument used for $F_d(s)$, one finds an absolute constant $c > 1$ so that

$$|G_d(s)| \ll c^{\nu(d)} |L(2s, \text{Sym}^2) \zeta(4s-2)^{-1}|,$$

for $\Re s > 3/4$ (say). Thus,

$$\begin{aligned} \sum_{n=1}^{\infty} \sum_{\substack{me^2=n \\ m_2d=r^2}} \frac{a(m)}{me} \prod_{p|4Nm_2d} \left(1 + \frac{1}{p}\right)^{-1} \exp(-n/X) \\ = \frac{1}{2} G_d(1)(\gamma + \log X) + \frac{1}{2} G'_d(1) + \mathbf{O}(c^{\nu(d)}), \end{aligned}$$

as $X \rightarrow \infty$. Inserting the main term into our sum, we obtain

$$\begin{aligned} \frac{6Y}{\pi^2 \phi(4N)} \sum_{\substack{\delta \leq X^{1/5} \\ (\delta, 4N)=1}} \frac{1}{\delta^2} \sum_{d|\delta^2} \frac{\tilde{\mu}(d)}{d} \\ \times \left\{ \frac{1}{2} \left(\gamma + \log \frac{Xd}{\delta^4} \right) \sum_{m_2d=r^2} \frac{a(m)}{m} \prod_{p|4Nm_2d} \left(1 + \frac{1}{p}\right)^{-1} \right. \\ \left. - \frac{1}{2} \sum_{m_2d=r^2} \frac{a(m) \log m}{m} \prod_{p|4Nm_2d} \left(1 + \frac{1}{p}\right)^{-1} \right\}. \end{aligned}$$

The second term gives an amount of $\mathbf{O}(Y)$. The contribution from the error term is also $\mathbf{O}(Y)$. The first term is

$$\frac{3Y}{\pi^2 \phi(4N)} (\log X) \sum_{\substack{\delta \leq X^{1/5} \\ (\delta, 4N)=1}} \frac{1}{\delta^2} \sum_{d|\delta^2} \frac{\tilde{\mu}(d)}{d} \sum_{m_2d=r^2} \frac{a(m)}{m} \prod_{p|4Nr} \left(1 + \frac{1}{p}\right)^{-1} + \mathbf{O}(Y).$$

Now using the definition of $\tilde{\mu}$, and making use of the fact that

$$\sum_{n=1}^{\infty} \frac{a(n)}{n^s} \left(\frac{\delta^2}{n} \right) = L(s) \sum_{d|\delta^2} \frac{\tilde{\mu}(d)}{d^s},$$

we obtain the identity

$$a(n) \left(\frac{\delta^2}{n} \right) = \sum_{\substack{d|\delta^2 \\ dm=n}} \tilde{\mu}(d) a(m).$$

Therefore, we find that the main term is

$$\frac{3Y(\log X)}{\pi^2 \phi(4N)} \sum_{\substack{\delta \leq X^{1/5} \\ (\delta, 4N)=1}} \frac{1}{\delta^2} \sum_{m_1} \frac{a(m_1)}{m_1} \sum_{\substack{r=1 \\ (r, 2N)=1}}^{\infty} \frac{a(r^2)}{r^2} \left(\frac{\delta^2}{r^2} \right) \prod_{p|4Nr} \left(1 + \frac{1}{p} \right)^{-1}.$$

In order to simplify this sum, consider the Dirichlet series

$$D_{\delta}(s) = \sum_{\substack{r=1 \\ (r, \delta)=1}}^{\infty} \frac{a(r^2)}{r^s} \prod_{p|4Nr} \left(1 + \frac{1}{p} \right)^{-1}.$$

For $\text{Re}(s) > 2$ it is an absolutely convergent series. Writing each r as $r = bu$ where $p|b \Rightarrow p|4N$ and $(u, 4N) = 1$, we see that

$$D_{\delta}(s) = \prod_{p|4N} \left(1 + \frac{1}{p} \right)^{-1} \sum_{\substack{b \\ p|b \Rightarrow p|4N}} \frac{a(b^2)}{b^s} \sum_{\substack{u=1 \\ (u, 4N\delta)=1}}^{\infty} \frac{a(u^2)}{u^s} \prod_{p|u} \left(1 + \frac{1}{p} \right)^{-1}.$$

As in Section 1, write

$$B_p(s) = \sum_{j=0}^{\infty} \frac{a(p^{2j})}{p^{js}}.$$

Then the sum over u is equal to

$$\prod_{p \nmid 4N\delta} \left(1 + \left(1 + \frac{1}{p} \right)^{-1} (B_p(s) - 1) \right)$$

and, factoring out B_p , we deduce that

$$D_{\delta}(s) = \prod_{p|4N} \left(1 + \frac{1}{p} \right)^{-1} \left(\prod_{p \nmid \delta} B_p(s) \right) \prod_{p \nmid 4N\delta} \left(1 - \frac{1}{p+1} \left(1 - \frac{1}{B_p(s)} \right) \right).$$

Notice that the last product above converges absolutely for $\text{Re}(s) > 1$ and at $s = 2$ can be bounded independently of δ . As for the second factor, we note

that it is

$$\left(\sum_{n=1}^{\infty} \frac{a(n^2)}{n^s} \right) \prod_{p|\delta} B_p(s)^{-1}$$

and at $s = 2$ we have (on using the estimate $a(p^{2j}) \leq (2j + 1)p^j$) that it is $\ll \delta/\phi(\delta)$.

Inserting this information into our sum, we see that it is

$$\frac{3Y(\log X)}{\pi^2 \phi(4N)} \sum_{\substack{\delta \leq X^{1/5} \\ (\delta, 4N)=1}} \frac{1}{\delta^2} D_{2N\delta}(2) \sum_{n_1} \frac{a(n_1)}{n_1}$$

which is equal to

$$\begin{aligned} & \frac{3Y(\log X)}{\pi^2 \phi(4N)} \prod_{p|4N} \left(1 + \frac{1}{p} \right)^{-1} \sum_{n_1} \frac{a(n_1)}{n_1} \left(\sum_{\substack{n_2=1 \\ (n_2, 2N)=1}}^{\infty} \frac{a(n_2^2)}{n_2^2} \right) \\ & \times \prod_{p \nmid 4N} \left(1 - \frac{1}{p+1} \left(1 - \frac{1}{B_p(2)} \right) \right) \\ & \times \sum_{\substack{\delta \leq X^{1/5} \\ (\delta, 4N)=1}} \frac{1}{\delta^2} \left(\prod_{p|\delta} B_p(2)^{-1} \right) \prod_{p|\delta} \left(1 - \frac{1}{p+1} \left(1 - \frac{1}{B_p(2)} \right) \right)^{-1}. \end{aligned}$$

By the estimate stated at the end of the previous paragraph, we may extend the sum over δ to infinity, thereby introducing an error of only $\mathbf{O}(YX^{-1/5+\varepsilon})$. Now let us simplify the sum over δ . As each summand is multiplicative in δ , we see that

$$\begin{aligned} & \prod_{p \nmid 4N} \left(1 + \frac{1}{B_p(2)} \left(1 - \frac{1}{p+1} \left(1 - \frac{1}{B_p(2)} \right) \right)^{-1} \left(\frac{1}{p^2} + \frac{1}{p^4} + \cdots \right) \right) \\ & = \prod_{p \nmid 4N} \left(1 + \frac{1}{B_p(2)(p^2-1)} \left(1 - \frac{1}{p+1} \left(1 - \frac{1}{B_p(2)} \right) \right)^{-1} \right). \end{aligned}$$

Multiplying this by

$$\prod_{p \nmid 4N} \left(1 - \frac{1}{p+1} \left(1 - \frac{1}{B_p(2)} \right) \right),$$

we find that it becomes

$$\prod_{p \nmid 4N} \left(1 - \frac{1}{p^2}\right)^{-1} \left(1 + \frac{1}{p} \left(\frac{1}{B_p(2)} - 1\right)\right).$$

We insert this calculation and the sum becomes

$$\begin{aligned} & \frac{1}{2} \frac{1}{\phi(4N)} (Y \log X) \sum_{n_1} \frac{a(n_1)}{n_1} \left(\sum_{(n_2, 2N)=1} \frac{a(n_2^2)}{n_2^2} \right) \\ & \times \prod_{p \nmid 4N} \left(1 - \frac{1}{p}\right) \prod_{p \nmid 4N} \left(1 + \frac{1}{p} \left(\frac{1}{B_p(2)} - 1\right)\right). \end{aligned}$$

In the notation of Remark 3 of Section 1,

$$\mathscr{P}(1) = \prod_{p \nmid 4N} \left(1 + \frac{1}{p} \left(\frac{1}{B_p(2)} - 1\right)\right).$$

The calculation in this remark shows that our sum is

$$\frac{1}{2} \frac{Y(\log X)}{\phi(4N)} \sum_{n_1} \frac{a(n_1)}{n_1} \sum_{\substack{n_2=1 \\ (n_2, 2N)=1}} \frac{a(n_2^2)}{n_2^2} \prod_{p \nmid 4Nn_2} \left(1 - \frac{1}{p}\right).$$

Summarizing, we have proved that the sum is

$$\frac{1}{4} CY(\log X) + O(Y) + O(X^{1/2} Y^{1/2} \log X \log \log X).$$

All of our estimates were made under the assumption that

$$Y \log^{-B} Y \leq X \leq Y(\log Y)^{1+\nu}$$

where $0 < \nu < \rho$. Combining the estimates of the sum and the integral, we deduce that if $X = Y/(\log Y)^\lambda$ for a small $\lambda > 0$, then the sum of the sum and the integral is

$$\begin{aligned} & \left(\frac{1}{\phi(4N)} \sum_{n_1} \frac{a(n_1)}{n_1} \sum_{n_2} \frac{a(n_2^2)}{n_2^2} \frac{\phi(4Nn_2)}{4Nn_2} \right) \left(\frac{1}{2} Y \log X - \frac{1}{2} Y \log \frac{X}{Y^2} \right) \\ & + O(Y(\log X)^{1-\nu}) \end{aligned}$$

and this is

$$\frac{1}{2} CY(\log Y) + O(Y(\log Y)^{1-\nu}),$$

which completes the proof of the theorem.

6. Concluding remarks

Our theorems can be generalised to include the case when D ranges over an arithmetic progression mod M . This gives rise to a more complicated constant. Though we have not developed it here, the method allows one to obtain asymptotic formulas of the form

$$\sum_{0 < -D \leq Y} L_D^{(j)}(1) = C_j Y \log^j Y + O(Y \log^{j-1} Y)$$

as $Y \rightarrow \infty$ for every $j \geq 1$. The results of these investigations will appear in a future work.

Acknowledgements. We would like to thank David Rohrlich, Henryk Iwaniec and the referee for their comments on a preliminary version of this paper.

McGILL UNIVERSITY, MONTREAL, CANADA
UNIVERSITY OF TORONTO, TORONTO, CANADA

REFERENCES

- [1] D. BUMP, S. FRIEDBERG, and J. HOFFSTEIN, Non-vanishing theorems for L -functions of modular forms and their derivatives, preprint.
- [2] A. S. FAINLEJB and O. SAPARNIJAZOV, Dispersion of real character sums and the moments of $L(1, \chi)$ (Russian), *Izv. Akad. Nauk. USSR, Ser. fiz.-mat. Nauk.* **19** Nr. 6 (1975), 24–29.
- [3] M. JUTILA, On the mean value of $L(1/2, \chi)$ for real characters, *Analysis* **1** (1981), 149–161.
- [4] V. A. KOLYVAGIN, Finiteness of $E(\mathbb{Q})$ and $\text{III}_{E/\mathbb{Q}}$ for a subclass of Weil curves (Russian), *Izv. Akad. Nauk. USSR, ser. Matem.* **52** (1988).
- [5] M. RAM MURTY, Oscillations of Fourier coefficients of modular forms, *Math. Ann.* **262** (1983), 431–446.
- [6] ———, Simple zeroes of certain L -series, *Proc. Number Theory Conf.* (ed. R. Mollin), Banff, Canada, 1987, pp. 427–439.
- [7] M. RAM MURTY and V. KUMAR MURTY, A variant of the Bombieri-Vinogradov theorem, in: *Number Theory*, ed. H. Kisilevsky and J. Labute, CMS Conf. Proc., **7** (1987), 243–272.
- [8] V. KUMAR MURTY, On the Sato-Tate conjecture, in: *Number Theory Related to Fermat's Last Theorem*, ed. N. Koblitz, Birkhauser-Verlag, Boston, 1982, pp. 195–205.
- [9] ———, Non-vanishing theorems for L -functions and their derivatives, *Automorphic Forms and Number Theory*, CRM, Montreal (1990), pp. 89–113.
- [10] R. RANKIN, Sums of powers of cusp form coefficients II, *Math. Ann.* **272** (1985), 593–600.
- [11] K. RUBIN, Tate-Shafarevic groups and L -functions of elliptic curves with complex multiplication, *Inv. Math.* **89** (1987), 527–559.
- [12] F. SHAHIDI, On certain L -functions, *Amer. J. Math.* **103** (1981), 297–356.

(Received April 24, 1989)

(Revised March 25, 1990)